

Case Report

Krackow's Technique Used to Repair for Two Month Old Bilateral Tendoachilles Rupture after Local Steroid Infiltration: Rare Case Report

Dasarath Kisan, Prafulla Kumar Sahoo, Hatia Marandi, Ranajit Panigrahi, Dibya Singha Das,
Satya Ranjan Patra, M Manoj

Orthopedic Department, Utkal University, Bhubaneswar, India.

Corresponding Author: Prafulla Kumar Sahoo

Received: 23/09/2015

Revised: 16/10/2015

Accepted: 23/10/2015

ABSTRACT

The causes for the painful heel around the Achilles tendon are Achilles tendinitis and retrocalcaneal bursitis.

The treatment modalities range from simple NSAID's, hot water fomentation, ultrasound phonophoresis, and intralesional steroids to surgery. There is a debate on giving intralesional steroids in and around the Achilles tendon. Most common complication of intralesional corticosteroids injection is rupture of the tendon. Here is a case of bilateral tendoachilles rupture due to intralesional corticosteroid infiltration. A 32 year old male suffering from bilateral retro calcaneal bursitis since two year was treated with two doses of intralesional steroid for left tendo-Achilles tendinitis 2 month apart and one dose intra lesional corticosteroid around tendoachilles on right Achilles tendon after 2 month of first dose of left side three months prior to rupture. Two months back patient had rupture of bilateral tendoachilles. This was confirmed by ultrasound and MRI examination. We treated the rupture using Krackow's technique to repair the tendon. Follow up at one year revealed good functional outcome. Bilateral tendoachilles rupture due to corticosteroid infiltration is very rare. Better to avoid intralesional steroid injections in and around the tendoachilles. MRI and ultrasound are confirmatory. MRI is more sensitive than ultrasound, but due to cost USG is the preferred investigation. Histopathological investigation reveals the tendon end to be degenerated; thereby chronic ruptures should be repaired surgically, which reduces chance of failure and improves functional outcome. In our case is operated in one step.

Key words: Bilateral tendoachilles rupture, Krackow's technique, MRI, Steroid infiltration, Ultrasound.

INTRODUCTION

The causes for the painful heel around the Achilles tendon are Achilles tendinitis and retro calcaneal bursitis. The treatment modalities range from simple NSAID's, hot water fomentation, ultrasonic therapy and intra lesional steroids to surgery. There is a debate on giving intralesional steroids in and around the Achilles tendon. Most common complication of intralesional

corticosteroids injection is rupture of the tendon. Here is a case of bilateral tendo achilles rupture due to intra-lesional corticosteroid infiltration.

CASE REPORT

A 32 yrs male working as a worker in tailoring shop, suffering from bilateral heel pain since two years, was treated with analgesics, physiotherapy and hot water fomentation. 9 months back patient was

diagnosed to have bilateral retro -calcaneal bursitis and treated with two doses of intra-lesional corticosteroid (Triamcinalone40 mg) on left side at an interval of 2 month. Patient was pain free after the injection. Two months ago while walking on a flat surface , patient felt a sudden snap in right ankle associated with

pain and discoloration of skin with swelling behind the ankle. Seven days later he had a similar episode in the left side. Since then the patient is unable to squat, stand for a few seconds without support, walk without support and unable to climb stairs.



Fig.1: Subcutaneous fat was absent, visible depression proximal to the insertion of tendoachilles



Fig. 2: No tenderness, palpable gap in the continuity of the tendoachilles proximal to its insertion, thickening noted proximal to the gap, firm in consistency



Fig. 3: No plantar flexion on squeezing the gastrosoleus muscles, indicates that discontinuity in tendoachilles



Fig. 4: Needle placed 10 cm proximal to the insertion of the tendoachilles does not move on plantar and dorsiflexion



Fig. 5: ankle joints show loss of normal regular configuration of kager's triangular space, which is between the superior aspect of the calcaneum and anterior aspect of tendoachilles



Fig. 6: USG using 12 mega hertz probe showed a heterogeneous hypo echoic area with discontinuity



Fig.7: MRI confirmed disruption of the tendon on T1 weighted and intratendinous generalized increased signal intensity on T2 weighted




Fig. 8: Krackow's technique with ethibond as the suture material

Examination: On inspection depigmentation was noted proximal to the insertion of tendoachilles (4cm x 4cm) Subcutaneous fat was absent, visible depression proximal to the insertion of tendoachilles (Fig. 1). No tenderness, palpable gap in the continuity of the tendoachilles proximal to its insertion, thickening not edproximal to the gap, firm in consistency (Fig. 2). Dorsiflexion - active 25°, passive 30°. Plantar flexion - active 20° and passive 30°. Thomsons test positive bilaterally (no plantar flexion on squeezing the gastro -soleus muscles, indicates that discontinuity intendoachilles) (Fig. 3). Obrien's needle test: positive (needle placed 10 cm proximal to the insertion of the tendoachilles does not move on plantar and dorsiflexion) (Fig. 4). Laboratory Investigations revealed patient is not suffering from diabetes mellitus. X- ray examination of the ankle joint show loss of normal regular configuration of kager's triangular space, which is between the superior aspect of the calcaneum and anterior aspect of tendoachilles (Fig. 5). USG using 12 mega hertz probe showed a heterogeneous hypo echoic area with discontinuity (Fig.6). MRI confirmed disruption of the tendon on T1 weighted and intra-tendinous generalized increased signal intensity on T2 weighted (Fig. 7).

We operated on both the ankles in one sitting. We used Krackow's technique with ethibond as the suture material (Fig. 8). Post operatively above knee cast was

applied for 2 weeks in equinus, suture removal done on 14th day. Short leg cast for next 2 weeks. After that, gradually foot is brought to plantigrade position and partial weight bearing started. At 6-8weeks full weight bearing over plantigrade short leg cast started. At 12 weeks reverse plantigrade ankle stop brace applied to gain full range of motion. Finally patient achieved 90% of the ankle movements at 6months and patient was able to walk and climb stairs without any discomfort.

DISCUSSION

The first description about the tendoachilles was given by Hippocrates. [1] The name Achilles tendon derived from the warrior of Homer's Iliad, Achilles. It is the strongest and thickest tendon in the body. [2] Thetis, mother of Achilles made his body invulnerable to physical harm by immersing him in the river Styx after learning of a prophecy that Achilles would die in the battle. However, the heel by which he was held remained untouched by the water and thus Achilles had a vulnerable point and was killed in a war, by a poisoned arrow fired in to his heel by the prince of Trojan. [3]

Bilateral Achilles tendon ruptures due to corticosteroid infiltration is rare. Spontaneous rupture most often occurs around 30 - 50 years, non-athlete sand women over the age of 50. Unilateral rupture of the Achilles tendon increases incidence of contralateral tendon rupture by more than 18 times. [4] Spontaneous

rupture may be due to congenital abnormal collagen, infection, rheumatism, endocrinal, neurological dysfunction, increasing age causing ischemic changes in Achilles tendon, excessive exercise leading to Achilles tendon degeneration intra-lesional/ oral steroids, norfloxacin, fluoroquinolones, high temperature and tendon calcification. [5] Normal tendoachilles consist of mainly type-1collagen, while a ruptured tendoachilles consists of type-3 collagen, which is less resistant to tensile forces and may, therefore, predispose the tendon to spontaneous rupture. [6] Patients will present with a sudden snap followed by difficulty in walking, gradually appearing swelling and discolouration of heel. Swelling at the site is due to internal bleeding which masks the actual discontinuity of tendon. The plantar tendon and flexor hallucis longus tendon provide sufficient plantar flexion strength so as to compensate for walking. [7] Examination reveals depigmentation of the skin, palpable gap, Thompsons test and Obrien's needle test will be positive. X ray examination is done to rule out any associated avulsion fractures of the calcaneum and shows loss of normal regular configuration of Kager's triangular space. [8] Real time high resolution Ultrasonography of Achilles tendon is more sensitive than soft tissue radiography. High frequency probes of 7.5 to 10.0megahertz provide the best resolution due to short focus and give dynamic and panoramic image of the tendon. It reveals heterogeneous hypo echoic areas with discontinuity. Spaces are filled with fluid. [9] Magnetic resonance imaging reveals disruption of the tendon on T1- weighted image and intra tendinous generalized increased signal intensity on T2 weighted image. [10] Bilateral chronic tendoachilles rupture can be treated both conservatively and surgically. The disadvantage of conservative treatment is a high re rupture rate. Amendala et al

reported a rerupture rate of 21% in conservative and 2% in operative cases. [11] Surgical repair of the tendon has different techniques like Krackow's, Lindholm, Lynn and Teuffer. In our case Krackow's technique has been used for the repair of the tendon. [12] Post operatively above knee cast immobilization done for 2 weeks. Suture removal done on the 14th day. Short leg cast with foot in equines is put for next 2 weeks. At 4 weeks, the cast is changed and foot is gradually brought to the plantigrade position, partial weight bearing is started over the next 2 weeks. At 6-8 weeks, short leg cast in plantigrade position is applied and full weight bearing started. At 12 weeks, a reverse 90 degree ankle stop brace is worn until the near normal range of motion is achieved and 80% strength. [13] In our case at 12 months patient has achieved full range of movements at both ankle joints and able to walk and climb stairs comfortably.

CONCLUSION

Bilateral tendoachilles rupture due to corticosteroid infiltration is very rare. Better to avoid intralesional steroid injections in and around the tendoachilles. MRI and ultrasound are confirmatory. MRI is more sensitive than ultrasound, but due to cost USG is the preferred investigation. Histopathological investigation reveals the tendon end to be degenerated; thereby chronic ruptures should be repaired surgically, which reduces chance of failure and improves functional outcome.

REFERENCES

1. Carden DG, Noble J, Chalmers J, Lunn P, Ellis J; Rupture of the calcaneal tendon. The early and late management. *Journal of Bone and Joint Surgery*, 1987; 69-B (3): 416-420.
2. O'Brien M; Functional anatomy and physiology often done; *Clinics in Sports Medicine*. 1992; 11(3): 505-520.

3. Shampo MA, Kyle RA; Medical mythology. Achilles. Mayo Clinic Proceedings, 1992; 67(7): 651.
4. Maffulli N, Irwin AS, Kenward MG, Smith F, Porter RW; Achilles tendon rupture and sciatica: a possible correlation. British Journal of Sports Medicine, 1998; 32(2): 174-177.
5. Unverferth LJ, Olix ML; The effect of local steroid injections on tendon. In Proceedings of the American Academy of Orthopaedic Surgeons. Journal of Bone and Joint Surgery, 1973; 55(A): 1315.
6. Coombs RRH, Klenerman L, Narcisi P, Nichols A, Pope EM; Collagen typing in Achilles tendon rupture. In Proceedings of the British Orthopaedic Research Society. Journal of Bone and Joint Surgery, 1980; 62-B (2): 258.
7. Hatstrup SJ, Johnson KA; A review of ruptures of the Achilles tendon. Foot Ankle, 1985; 6(1): 34-38.
8. Arner O, Lindholm Å, Lindvall N; Roentgen changes in subcutaneous rupture of the Achilles tendon. ActaChirurgica Scandinavica, 1959; 116(5-6): 496-500.
9. Fornage BD, Rifkin MD; Ultrasound examination of tendons. Radiologic Clinic of North America, 1988; 26(1): 87-107.
10. Kabbani YM, Mayer DP; Magnetic resonance imaging of tendon pathology about the foot and ankle. Part I. Achilles tendon. Journal of American Pediatric Medical Association, 1993; 83(7): 418-420.
11. Willits K, Amendola A, Bryant D, Mohtadi NG, GiffinJR, Fowler P *et al.*; Operative versus non operative treatment of acute achilles tendon ruptures: A Multicenter Randomized Trial Using Accelerated Functional Rehabilitation. J. Bone Joint Surg Am., 2010; 92(17): 2767-2775.
12. Krackow KA, Thomas SC, Jones LC; A new stitch for ligamentous tendon fixation: brief note. J Bone Joint Surg Am., 1986; 68(5): 764-766.
13. Kangas J, Pajala A, Ohtonen P, Leppilahti J; Achilles tendon elongation after rupture repair: a randomized comparison of 2 postoperative regimens. Am J Sports Med., 2007; 35(1): 59-64.

How to cite this article: Kisan D, Sahoo PK, Marandi H et al. Krackow's technique used to repair for two month old bilateral tendoachilles rupture after local steroid infiltration: rare case report. Int J Health Sci Res. 2015; 5(11):451-456.
