

Original Research Article

## Relationship between Body Mass Index and Blood Pressure Level in Hypertensive Patients Attending Kiambu District Hospital, Kenya

Jane Mbijiwe<sup>1</sup>, Peter Chege<sup>2</sup>, Ann Munyaka<sup>2</sup>

<sup>1</sup>Master of Science (Food Nutrition and Dietetics) Student, <sup>2</sup>Lecturer,  
Department of Nutrition and Dietetics, Kenyatta University, Kenya, P.O. BOX 43844, Nairobi, Kenya

Corresponding Author: Jane Mbijiwe

### ABSTRACT

Uncontrolled hypertension is the primary cause of cardiovascular disease occurrence which greatly contributes to fatality worldwide. Obesity is known to impact cardiovascular health, being an appropriate predictor of blood pressure. Many hypertensive patients have been found to be obese across the world. Scarce information exists on the relationship between obesity and blood pressure level among hypertensive patients. This research aimed at assessing the body mass index of hypertensive patients in a developing country. The study was conducted on a sample of 134 hypertensive patients attending Kiambu District Hospital in Kenya. Cross-sectional analytical research design was adopted and respondents selected using systematic sampling method. Anthropometric parameters of weight and height were used to assess body mass index. A pretested questionnaire was used to collect data from the respondents. Data was analyzed by use of statistical package for social science. The respondent's body mass index was described by use of descriptive statistics. The relationship between blood pressure level and body mass index was determined by use of Pearson correlation test. A p-value of <0.05 was used as criterion for statistical significance. The study population had high body mass index as revealed by the high prevalence of overweight and obesity at 82.9%. Of the 134 respondents, 79.1% had uncontrolled blood pressure. Positive significant relationship was found between body mass index and blood pressure ( $r=0.683$ ,  $p<0.001$ ). Based on these findings, measures that would lead to healthy body mass index among hypertensive patients should be put in place for better hypertension management.

**Key Words:** Body mass index, hypertension, overweight, obesity

### INTRODUCTION

Hypertension accounts for 10% of health care spending globally. [1] It is a modifiable risk factor for renal disease and cardiovascular disease. Hypertension is defined as a systolic blood greater than 140 mm Hg and a diastolic blood pressure greater than 90 mm Hg based on the average of two or more correct blood pressure measurement taken by a health care provider. Among hypertensive patients worldwide, only 31% have optimal blood pressure levels. [2] The Kenya Stepwise Survey Report [3] indicated that the

prevalence of uncontrolled hypertension among hypertensive patients in Kenya is at 24%. Many hypertensive patients regularly attend hospital due to health complications caused by the disease. In Kiambu County, a rise in hypertension prevalence has been observed. The disease is ranked seventh and fifth in causing fatality and hospital admission respectively in the County. [4]

Uncontrolled hypertension damages cells lining coronary vessels resulting in inflammation, vascular weakness and scarring, blood clots, blocked arteries and plaque formation. [5-8] Prolonged periods of

uncontrolled hypertension affect heart functions by weakening heart muscles and enlarging the heart which may lead to death. Glomerulosclerosis is caused by uncontrolled hypertension when blood vessels are damaged by high blood pressure. [9] The most appropriate predictor of blood pressure level has been reported to be nutrition status hence important for maintenance of optimal blood pressure levels. Obesity is known to impact cardiovascular health globally and is related to uncontrolled hypertension. Obesity is associated with 16% stroke related deaths. [10] Studies have indicated that the risk of developing hypertension independent of physical inactivity and unhealthy diets is increased 3 times in overweight people. [11-12] An estimated 27% people in Kenya are overweight or obese. [13] Obesity affects the rennin-angiotensin aldosterone system increasing absorption of renal sodium which results in elevated blood pressure. Obesity is also known to cause insulin resistance leading to sodium retention. Obesity increases the risk of heart failure since excess weight raises cardiac output lowering peripheral resistance. Weight reduction through increased physical activity and dietary changes help in maintenance of optimal blood pressure. [14] It is important to sensitize hypertensive patients on the benefits of behaviour change and put in place measures that encourage adoption of healthy practices.

Several anthropometric measures have been identified as measures of obesity with BMI being found the most appropriate due to its accuracy and reproducibility. [15] This study therefore used this parameter to determine nutrition status. Achieving and maintenance of optimal blood pressure levels in hypertensive patients is beneficial for the prevention of hypertension complications. There is paucity of published research findings on the relationship between blood pressure and BMI of hypertensive outpatients in Kenya hence the need for this study.

Increased hypertension prevalence in health populations has been observed with increased age. [16] Advanced age impairs the function of the heart by altering the vascular structure leading to arterial stiffness that decreases buffering hence increasing blood pressure. Studies have reported that the age group with the highest risk of uncontrolled blood pressure among hypertensive patients is 18-57 years. [16-17]

## METHODS

A Cross-sectional analytical research design was adopted in this study. The study was carried out at Kiambu District Hospital in Kiambu County, Kenya. The target population was hypertensive out-patients aged between 18-57 years attending the outpatient clinic at Kiambu District Hospital. The identified patients were on regular medical check-ups and follow up at the hospital based on their blood pressure levels. Hypertensive patients attending the hypertensive clinic for the first time, expectant women and those with health complications including diabetes, kidney and heart diseases were excluded from the study. Participants were identified by use of systematic sampling technique. Data collection was carried out for three months to obtain the sample size of 134.

## RESEARCH INSTRUMENTS

Anthropometry data was collected by use of anthropometric parameters of height and weight followed by calculation of BMI. The respondents stood straight with minimum clothing with no shoes for the height to be taken using a stadiometer and a wooden head board. The height was measured to the nearest 0.5 centimetres. A weighing machine calibrated to zero was used to measure weight recorded to the nearest 0.05 kilogram. The measurements were taken thrice and the average calculated. BMI was determined by dividing weight in (kg) by height squared ( $m^2$ ) BMI was then categorized according to WHO body mass index cut off points of underweight (BMI<18.5), normal weight

(BMI; 18.5–24.9), overweight (25–29.9) and obesity (BMI>30).

Mercury sphygmomanometers were used to take blood pressure measurements. The respondents were allowed five minutes relaxation before the test was conducted. The client was seated with legs uncrossed back and arm supported. The reading was done at the cuff where all clothing around the area was loose. Three readings were taken with a repeat being done at 3-5 minutes interval and an average calculated and blood pressure levels then categorized using WHO cut off points.

## DATA ANALYSIS AND PRESENTATION

Before data entry the filled questionnaires were coded, checked and cleaned for consistency. The data was then entered into the computer for analysis. Statistical package for the social science was used to analyse data from blood pressure level and anthropometry. BMI levels of the study population were described by use of descriptive statistics. Pearson correlation coefficient test established the relationship between blood pressure level and BMIA p-value of <0.05 was used as a criterion for statistical significance.

### Ethical Clearance

Approval was sought from Kenyatta University Graduate School. Ethical clearance was obtained from Kenyatta University Ethical Review Committee. A research permit was obtained from the National Commission for Science, Technology and Innovation (NACOSTI). Permission was also sought from Kiambu Hospital management to access the medical records of the patients. Respondents participated in the research based upon their informed consent and assurance of confidentiality by the researcher.

## RESULTS

### Hypertensive characteristics

Majorities (69.4%) of the respondents were women and only 30.6% were men. The age group with the largest number of respondents (38.8%) was 50-57 years old. Overall 76.1% of the respondents had been diagnosed with hypertension in less than 5 years with majority (64.2%) visiting the hypertensive clinic for the second time. Approximately 79.1% of respondents had a SBP equal to or greater than 140 mmHg while 72.4% had a DBP equal to or greater than 90 mmHg. The mean Systolic blood pressure was  $146.23 \pm 0.87$  and the diastolic blood pressure  $92.06 \pm 0.12$ .

Hypertensive Characteristics						
Variable	Male		Female		Total	
	n	%	n	%	n	%
Age in Years						
18-25	6	14.6	14	15.1	20	14.9
26-33	5	12.2	9	9.7	14	10.4
34-41	5	12.2	20	21.5	25	18.7
42-49	10	24.4	13	14.0	23	17.2
50-57	15	36.6	37	39.8	52	38.8
Length of illness						
< 5 years	25	61.0	77	82.8	102	76.1
5-10 years	14	34.1	9	9.7	23	17.2
>10 years	2	4.9	7	7.5	9	6.7
No. of Clinic Visits						
2 times	27	65.9	59	63.4	86	64.2
3-4 times	8	19.5	19	20.4	27	20.1
>5 times	6	14.6	15	16.1	21	15.7
Systolic Blood pressure						
< 140 mmHg	16	39.0	12	12.9	28	20.9
>=140 mmHg	25	61.0	81	87.1	106	79.1
Diastolic Blood pressure						
< 90 mmHg	15	36.6	22	23.7	37	27.6
>=90 mmHg	26	63.4	71	76.3	97	72.4

Table 1: Respondents Demographic characteristics, health seeking behaviour and blood pressure levels

### Nutrition Status of the Respondents

Nutrition status of the study subjects was determined by use of WHO (2004) BMI cut off points. The average BMI of the study respondents was  $27.63 \pm 1.12$ . The study findings indicated that majority (59.0%) of the respondents were overweight, while 23.9% were obese.

When cross tabulation was done to compare BMI and age, the results showed that majority (33) of the overweight respondents were in the 50-57 years age group.

**Table 2: Cross tabulation BMI & Age**

BMI	Age in Years					Total
	18-25	26-33	34-41	42-49	50-57	
Underweight	2	0	1	0	9	12
Normal	6	0	2	1	2	11
Overweight	8	11	12	15	33	79
Obese	4	3	6	5	14	32

### Relationship between BMI and blood pressure

A positive significant relationship ( $r=0.772$ ,  $p=0.028$ ) was observed between BMI and SBP, this meant that SBP increased with increased BMI.

**Table 3: Relationship between BMI and blood pressure**

Variable	r	p-value
BMI and systolic blood pressure	0.772*	0.028
BMI and diastolic blood pressure	0.444	0.849

\*Correlation is significant at the 0.05 level (2-tailed).

### DISCUSSION

Age is both a risk factor for onset of hypertension and uncontrolled hypertension. In this study majority of the respondents were in the age group of 50-57 years this finding is in agreement with other studies that have reported blood pressure level increases with age. [18-20] In the current study age was found to influence BMI establishing a trend of increased weight with age, findings that are in agreement with other studies that have linked weight gain to age. [21-23] Approximately 79.1% of respondents had a SBP equal to greater than 140 mmHg while 72.4% had a DBP equal to or greater than 90 mmHg. These findings show that majority of the respondents in the current study had sub optimal blood pressure level.

In this study, 82.9% of the respondents were both overweight and obese. Similar findings of a majority of hypertensive patients being overweight and obese have been reported in other studies. [24-26] This finding could be explained by the possible fact that the respondents were consuming unhealthy diets and had low levels of physical activity. The current study findings showed a significant relationship between blood pressure and BMI. The relationship between hypertension and obesity has been documented globally with

[27-28] reporting significant relationship between obesity and hypertension.

### CONCLUSION

High prevalence of overweight and obesity, and uncontrolled hypertension was observed in the current study. This indicates the urgent need for hypertensive patient's behavior overhaul to increased physical activity engagement and healthy dietary intake for achievement and maintenance of healthy body weight. In this regard, health professionals should consider early nutrition screening of hypertensive patients to prevent overweight and obesity. Intervention programs that create awareness on the importance of health lifestyle for optimal blood pressure levels should be developed by relevant policy makers.

### ACKNOWLEDGEMENT

I acknowledge with appreciation my supervisors Dr. Peter Chege and Dr. Ann Munyaka both of Kenyatta University for their guidance, support and expertise contribution for the successful completion of this work. I also thank the hypertensive patients at Kiambu Hospital who participated in this study.

### REFERENCES

1. Campbell NR, Lackland DT, Niebylski ML, Nilsson PM. Is reducing dietary sodium controversial? Is it the conduct of studies with flawed research methods that is controversial? A perspective from the World Hypertension League Executive Committee. The Journal of Clinical Hypertension. 2015; 17(2):85-6.
2. Van de Vijver S, Akinyi H, Oti S, Olajide A, Agyemang C, Aboderin I, Kyobutungi C. Status report on hypertension in Africa-Consultative review for the 6th Session of the African Union Conference of Ministers of Health on NCD's. Pan African Medical Journal. 2014;16(1).
3. Ministry of Health Kenya. Non Communicable Diseases Risk Factors Report. Kenya: KNBS; 2015. 13p
4. Eston N, Samuel K, Paul S. Exploring Kenyas Inequality. Pulling Apart or Pooling together? Kenya: Ascent

- Limited; 2013. 119p.
5. Mutua EM, Gitonga MM, Mbuthia B, Muiruri N, Cheptum JJ, Maingi T. Level of blood pressure control among hypertensive patients on follow-up in a regional referral hospital in Central Kenya. *The Pan African medical journal*. 2014;18.
  6. Ferdinand KC. Cardiovascular disease in blacks: can we stop the clock? *The Journal of Clinical Hypertension*. 2008; 10(5):382-9.
  7. Radhika G, Sathya RM, Sudha V, Ganesan A, Mohan V. Dietary salt intake and hypertension in an urban south Indian population-[CURES-53]. *Journal of Association of Physicians of India*. 2007;55(6):405-11.
  8. Farah R, Zeidan RK, Chahine MN, Asmar R, Chahine R, Salameh P, Pathak A, Hosseini H. Predictors of Uncontrolled Blood Pressure in Treated Hypertensive Individuals: First Population-Based Study in Lebanon. *The Journal of Clinical Hypertension*. 2016;18(9):871-7.
  9. Low KJ, Pelter MA, Deamer RL, Burchette RJ. Identification and evaluation of risk factors in patients with continuously uncontrolled hypertension. *The Journal of Clinical Hypertension*. 2015;17(4):281-9.
  10. Kowal P, Chatterji S, Naidoo N, Biritwum R, Fan W, Lopez Ridaura R, Maximova T, Arokiasamy P, Phaswana-Mafuya N, Williams S, Snodgrass JJ. Data resource profile: the World Health Organization Study on global Ageing and adult health (SAGE). *International journal of epidemiology*. 2012;41(6): 1639-49.
  11. Narkiewicz K. Diagnosis and management of hypertension in obesity. *Obesity reviews*. 2006;7(2):155-62.
  12. World Health Organization. A global brief on hypertension: silent killer, global public health crisis: World Health Day 2013.
  13. Mathenge W, Foster A, Kuper H. Urbanization, ethnicity and cardiovascular risk in a population in transition in Nakuru, Kenya: a population-based survey. *BMC public health*. 2010;10(1):569.
  14. El Banna S, Fouad A. Prevalence of pre-hypertension and hypertension in a sample of Egyptian adults and its relation to obesity. *Australian Journal of Basic and Applied Sciences*. 2012;6 (13):481-9.
  15. Feng RN, Zhao C, Wang C, Niu YC, Li K, Guo FC, Li ST, Sun CH, Li Y. BMI is strongly associated with hypertension, and waist circumference is strongly associated with type 2 diabetes and dyslipidemia, in northern Chinese adults. *Journal of epidemiology*. 2012; 22(4):317-23.
  16. Cohen L, Curhan GC, Forman JP. Influence of age on the association between lifestyle factors and risk of hypertension. *Journal of the American Society of Hypertension*. 2012;6(4): 284-90.
  17. Svetkey LP, Erlinger TP, Vollmer WM, Feldstein A, Cooper LS, Appel LJ, Ard JD, Elmer PJ, Harsha D, Stevens VJ. Effect of lifestyle modifications on blood pressure by race, sex, hypertension status, and age. *Journal of human hypertension*. 2005;19(1):21.
  18. Babiker FA, Elkhalfifa LA, Moukhyer ME. Awareness of hypertension and factors associated with uncontrolled hypertension in Sudanese adults: cardiovascular topic. *Cardiovascular journal of Africa*. 2013;24(6):208-12.
  19. McEniery CM, Wilkinson IB, Avolio AP. Age, hypertension and arterial function. *Clinical and Experimental Pharmacology and Physiology*. 2007; 34(7):665-71.
  20. Rockwood MR, Howlett SE. Blood pressure in relation to age and frailty. *Canadian geriatrics journal: CGJ*. 2011; (1):2.
  21. Eshkoor SA, Hamid TA, Shahar S, Ng CK, Mun CY. Factors Affecting Hypertension among the Malaysian Elderly. *Journal of Cardiovascular Development and Disease*. 2016;3(1):8.
  22. Mungreiphy NK, Kapoor S, Sinha R. Association between BMI, blood pressure, and age: study among Tangkhul Naga tribal males of Northeast India. *Journal of Anthropology*. 2011;2011.
  23. Mártires MA, Costa MA, Santos CS. Obesity in aged hypertension patients.

- Texto & Contexto-Enfermagem. 2013; 22(3):797-803.
24. Batiha AM, AlAzzam M, ALBashtawy M, Tawalbeh L, Tubaishat A, Alhalaiqa FN. The relationship between hypertension and anthropometric indices in a Jordanian population. *Advanced Studies in Biology*. 2015; 7(5):233-43.
25. Deji SA, Olayiwola IO, Fadupin GT. Assessment of Nutritional Status of a Group of Hypertensive Patients Attending Tertiary Healthcare Facilities in Nigeria. *East African medical journal*. 2014;91(3):99-104.
26. Motlagh SF, Chaman R, Sadeghi E, Eslami AA. Self-Care Behaviors and Related Factors in Hypertensive Patients. *Iranian Red Crescent Medical Journal*. 2016;18(6).
27. Asgedom SW, Gudina EK, Desse TA. Assessment of Blood Pressure Control among Hypertensive Patients in Southwest Ethiopia. *PloS one*. 2016; 11(11):e0166432.
28. Al-Wehedy A, Abd Elhameed SH, Abd El-Hameed D. Effect of lifestyle intervention program on controlling hypertension among older adults. *Journal of Education and Practice*. 2014;5(5):61

How to cite this article: Mbijiwe J, Chege P, Munyaka A. Relationship between body mass index and blood pressure level in hypertensive patients attending Kiambu district hospital, Kenya. *Int J Health Sci Res*. 2017; 7(10):224-229.

\*\*\*\*\*