

Original Research Article

Nutritional Factors and Vitamin D Status in Relation to Severity of Pneumonia among Children

Dr. Priya PS¹, Dr. Elizabeth KE²

¹Junior Resident, ²Professor & Head,
Dept. of Pediatrics, SAT Hospital, Govt. Medical College, Thiruvananthapuram, Kerala, India-695011

Corresponding Author: Dr. Elizabeth KE

ABSTRACT

Objective: To compare nutritional factors like Infant & Young Child Feeding (IYCF) practices, nutritional status and serum vitamin D levels with severity of pneumonia among 1-5-year-old children.

Methods: Children admitted with severe and very severe pneumonia, as per IMNCI guidelines were enrolled. Their nutritional status including IYCF practices and serum 25-hydroxy vitamin D levels were compared.

Results: Out of 78 children, 40 (51%) had severe and 38 (49%) had very severe pneumonia. Socio-demographic factors and nutritional status were comparable in the two groups ($P > 0.05$). Majority, 22/34 (65%) with non-exclusive breastfeeding for 6 months, 7/9 (78%) with duration of breastfeeding < 4 months of age and 26/37 (70%) among those with previous admission for pneumonia had very severe pneumonia ($P < 0.05$). Vitamin D was suboptimum or deficient in 37 (48%). 25/37 (68%) in this category had very severe pneumonia ($P < 0.01$); 10/11 in the deficient, 6/6 in the severely deficient and 3/3 in the very severely deficient group had very severe pneumonia ($P < 0.001$). 34/37 (92%) with previous history of pneumonia were in the deficient group ($P < 0.001$).

Conclusion: Very severe pneumonia was significantly more in those with non-exclusive breastfeeding for 6 months, duration of breastfeeding < 4 months of age, previous admission with pneumonia and vitamin D deficiency. Very severe pneumonia was more in those with more severe vitamin D deficiency. Nutritional aspects like IYCF practices and vitamin D status being modifiable factors, the results of the study are of utmost public health importance.

Key Words: Severity of Pneumonia, IYCF practices, Nutritional status, Vitamin D Deficiency

INTRODUCTION

Pneumonia is the most common killer disease among children. Nutritional factors like Infant & Young Child Feeding (IYCF) practices and nutritional status influence the occurrence and outcome in community acquired pneumonia (CAP). The anti-rachitic or sunshine factor, vitamin D is now considered as a steroid hormone that regulates genomic functions like regulation of cell growth, cell differentiation and immune function in addition to bone health. The extra skeletal effects and the universal

presence of vitamin D receptors have been highlighted. ⁽¹⁾ The change in lifestyle, dressing habits, lack of sun exposure and environmental pollution has resulted in higher prevalence of vitamin D deficiency. It has also been linked to respiratory infections such as pneumonia, tuberculosis, bronchiolitis and asthma. ⁽²⁾ This study was undertaken to assess nutritional factors and vitamin D status in relation to severity of pneumonia among children.

METHODS

Participants consisted of 78 children in the age group, 1-5 years of age; 40 with severe and 38 with very severe pneumonia. Sample size was calculated as 78 , based on the formula $(Z\alpha)^2 \times SD^2 / d^2$. Consecutive cases, with severe and very severe pneumonia, as per IMNCI guidelines, (3) admitted in a tertiary care teaching hospital were enrolled. Institutional Ethics Committee approval and informed consent from the parents were obtained prior to enrolment. Those on vitamin D supplements, having rickets or other co-morbidities like renal and liver diseases and those on anticonvulsant and antitubercular therapy were excluded from the study. Socio-demographic factors, IYCF practices, nutritional status and serum 25-hydroxy vitamin D levels were assessed. Dietary intake of vitamin D and extent of sun exposure were not assessed. Nutritional status was assessed using the WHO growth charts. (4) The 25-OH-Vitamin-D-ELISA, 96-welled kit, manufactured by Biovendor research and diagnostics Products, Hamburg, Germany (Cat No. REA 300/96), using monoclonal anti-25-OH Vitamin D antibody was used. Serum 25-hydroxy vitamin D levels were categorized as: > 30 – optimum/sufficiency, $20-30$ – suboptimum/insufficiency, $10-20$ – deficiency, $5-10$ – severe and < 5 ng/ml – very severe deficiency. (5)

Data collected using structured Performa was managed using MS Excel software and analyzed using SPSS version 17.0 for Windows. Descriptive statistics was used for participant characteristics and Fisher's exact and Chi square tests were used for proportions.

RESULTS

Out of the 78 children, 40 (51%) had severe and 38 (49%) had very severe pneumonia. The male to female ratio was 49%:51%. 43 (55%) were in the age group 1-3 years and the rest were 3-5 years old. Socio-demographic variables like age, gender, maternal education, maternal

employment, place of residence, housing conditions, place of residence and socio-economic status were comparable in the two groups with severe and very severe pneumonia ($P > 0.05$). All of them belonged to middle and lower class. Only 44 (56%) of them had exclusive breastfeeding till six months of age and 9 (11.5%) had breastfeeding only for 4 months. Majority, 22/ 34 (65%) with non-exclusive breastfeeding for 6 months ($P < 0.05$) and 7/9 (78%) with duration of breastfeeding < 4 months of age had very severe pneumonia ($P < 0.01$). Significant protective nature of exclusive breast feeding for six months against very severe pneumonia was noted ($P < 0.01$; $\chi^2 = 6.167$).

33 (42%) were underweight, 21 (27%) had stunting and 37 (46%) had wasting and 3 children (4%) were severely underweight with severe wasting and 1 (1.2%) had severe stunting. Nutritional status was comparable in the two groups ($P > 0.05$).

Table 1. Mean Vitamin D Level (SD) in the Two Subgroups

Category	No. (%)	Mean Vitamin D Level- ng/ml (SD)
Severe Pneumonia	40 (51)	36.02 (10.8)
Very Severe Pneumonia	38 (49)	22.83 (13.6)
Combined	78 (100)	29.6 (13.9)

Vitamin D was suboptimum or deficient in 37 (48%). The levels ranged between 2-70 ng/ml with a mean of 36.02 ± 10.8 in severe and 22.83 ± 13.6 ng/ml in the very severe group. The mean serum 25 hydroxy Vitamin D levels (SD) are given in Table 1. The median (Inter Quartile Range-IPR) vitamin D levels in the two subgroups are depicted in fig 1. 25/37 (68%) in the deficient/suboptimum category had very severe pneumonia ($P < 0.01$, $\chi^2 = 10.01$). Very severe pneumonia was more in those who had more vitamin D deficiency; 10/11 in deficient, 6/6 in the severely deficient and 3/3 in the very severely deficient group had very severe pneumonia ($P < 0.001$). None of the socio-demographic variables had any association with vitamin D deficiency (P

>0.05). Table 2 summarizes the proportion of children with vitamin D sufficiency and deficiency in the two subgroups.

37 (47%) gave a previous history of admission for pneumonia. 26/37 (70%) among those with previous admission for

pneumonia had very severe pneumonia (P < 0.05). Majority, 34/37 (92%) with previous history of pneumonia were in the vitamin D deficient group (P <0.001(P <0.0001; $\chi^2 = 55.79$).

Table 2. Proportion of Children with Vitamin D Sufficiency and Deficiency in the Two Subgroups.

Vitamin D Status	Severe Pneumonia	Very Severe Pneumonia	Total No.	Percentage
	No. (%)	No. (%)		
Optimum > 30ng/ml	28 (70)	13 (34.2)	41	52.6
Suboptimum 20-30ng/ml	11 (27.5)	6 (15.79)	17	21.79
Deficiency 10-20 ng/ml	1 (2.5)	10* (26.32)	11	14.1
Severe Deficiency 5-10 ng/ml	0	6* (15.79)	6	7.7
Very Severe Deficiency <5 ng/ml	0	3* (7.89)	3	3.8
Total	40 (100)	38 (100)	78	100

* P < 0.001 Significant




Fig. 1 Median (Inter Quartile Range) Vitamin D levels with minimum and maximum values in children with Severe and Very Severe Pneumonia

DISCUSSION

In the present study, socio-demographic variables like age, gender, maternal education, maternal employment, place of residence, housing conditions, place of residence and socio-economic status were comparable in those with severe and very severe pneumonia. All of them belonged to middle and lower class, representing the class of people attending the public sector for medical care. Factors like maternal education, family income and availability of medical care have been attributed as risk factors for CAP. (6) The Pneumonia Etiology Research for Child

Health (PERCH) Study had found gross variability in these factors depending up on the location of study. (7)

Nutritional status was comparable in the two groups and only three children had severe wasting and one had severe stunting. High mortality has been reported in those with severe malnutrition. (8) The state of Kerala has been acknowledged as a high performing state with steady decline in child malnutrition and mortality rates. (9) IYCF practices like non-exclusive breastfeeding for 6 months and duration of breastfeeding <4 months of age were associated with very severe pneumonia. The protective role of breastfeeding has been amply highlighted by earlier workers. (10) A study had reported that subclinical vitamin D deficiency and nonexclusive breastfeeding in the first 4 months of life were significant risk factors for severe acute lower respiratory infection in Indian children. (11)

Very severe pneumonia was more in the vitamin D deficient/suboptimum category and severity was more with more severe deficiency. These observations are in accordance with other reported studies. (11,12) A similar study reported a mean level of 34.8 ± 15.6 ng/ml in severe pneumonia and 19.6 ± 9.8 ng/ml in very severe group

compared to 36.02 ± 10.8 in severe and 22.83 ± 13.6 g/ml in the very severe group in the present study. ⁽¹³⁾ It has been shown that after adjusting for demographic factors, clinical data, and season, 25(OH)D levels <30 ng/ml were associated with 56% higher odds of CAP compared to levels ≥ 30 ng/ml ⁽¹⁴⁾ Significant increase in occurrence of severe CAP in children with vitamin D levels <20 ng/ml has been reported. ⁽¹⁵⁾ Various case control trials have shown positive impact of vitamin D supplementation on the outcome of children with respiratory diseases. ⁽¹⁶⁻¹⁸⁾

47% gave a history of previous admission for pneumonia, comparable to another study that showed previous history of pneumonia in 41%. ⁽¹⁹⁾ It was interesting to note that the proportion was strikingly more in the vitamin D deficient group, up to 92%. The study highlights that optimizing IYCF practices and addressing severe vitamin D deficiency in children can help to reduce morbidity and mortality in pneumonia, which is a major killer disease in children. ⁽²⁰⁾

CONCLUSION

Non-exclusive breastfeeding for 6 months, duration of breastfeeding <4 months of age, previous admission for pneumonia and vitamin D deficiency as well as severity of deficiency were related to severity of pneumonia, which is a major killer disease in children. Previous admission for pneumonia was also more in those with vitamin D deficiency. Nutritional aspects including vitamin D status being modifiable factors, the results are reported due to its public health importance.

ACKNOWLEDGEMENT

The financial support from SAT Endowment Scheme is gratefully acknowledged. This study had won Dr. Omana Mathew Endowment Award of Indian Academy of Pediatrics, Kerala State Branch for the best MD Pediatrics Thesis from the Kerala University for Health Sciences (KUHS) for the year 2015.

Contributors: PPS collected and analyzed the data. EKE conceived the idea, supervised the study and drafted the paper.

REFERENCES

1. Wacker M, Holick MF, Vitamin D-effects on skeletal and extraskeletal health and the need for supplementation. *Nutrients*. 2013; 5:111-148.
2. Herr C, Greulich T, Kocuzilla RA, Meyer S, Zakharkina T, Branscheidt M, Eschmann R, Bals R. The role of vitamin D in pulmonary disease: COPD, asthma, infection, and cancer. *Respiratory Research*. 2011; 12:31-33
3. Government of India, Ministry of Health and Family Welfare. IMNCI training module for physicians. Government of India, 2003.
4. WHO Growth Reference Data for 0-5 years. Available from <http://www.who.int/growthref/en/>.
5. Holick MF, Binkley NC, Bischoff-Ferrari A, Gordon CM, Hanley DA, Heaney RP, Murad MH, Weaver CM. Evaluation, Treatment, and Prevention of Vitamin D Deficiency: An Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab*. 2011; 96 (7): 1911-1930. DOI: <https://doi.org/10.1210/jc.2011-0385>
6. HameedAzab SFA, Sherief LM, Saleh SH, Elsaeed WF, Elshafie MA, Abdelsalam SM. Impact of the socioeconomic status on the severity and outcome of community-acquired pneumonia among Egyptian children: A cohort study. *Infectious Diseases of Poverty*. 2014; 3:14-17 DOI: 10.1186/2049-9957-3-14
7. Wonodi CB, Deloria-Knoll M, Feikin DR, DeLuca AN, Driscoll AJ, MoisiJC, Johnson HL, David R. Murdoch DR, O'Brien KL, Levine OS, Scott JAG and the Pneumonia Methods Working Group and PERCH Site Investigators. Evaluation of risk factors for severe pneumonia in children: The Pneumonia Etiology Research for Child Health Study. *Clin Infect Dis*. 2012 Apr 1; 54(Suppl 2): S124-S131. doi: 10.1093/cid/cir1067

8. Chisti MJ, Tebruegge M, La Vincente S, Graham SM, Duke T. Pneumonia in severely malnourished children in developing countries—mortality risk, aetiology and validity of WHO clinical signs: a systematic review. *Trop Med Int Health* 2009; 14: 1173–89.
9. National Family Health Survey- 4. Key findings from NFHS-4 fact sheets for key indicators based on final data-Kerala, 2015-16. rchiips.org/NFHS/factsheet_NFHS-4.shtml
10. Broor S, Pandey RM, Ghosh M, Maitreyi RS, Lodha R, Singhal T, Kabra SK. Riskfactors for Severe acute lower respiratory tract infection in Under-Five Children. *Indian Pediatr.* 2001; 38: 1361-1369
11. Wayse V, Yousafzai A, Mogale K, Filteau S. Association of subclinical vitamin D deficiency with severe ALRI in Indian children under 5 years; *Eur J Clin Nutr.* 2004; 58: 563-567
12. Remmelts HHF, van de Garde EMW, Meijvis SCA, Peelen ELGCA, Damoiseaux JGMC, Grutters JC, Biesma DH, Bos WJWB, Rijkers GT. Addition of vitamin D status to Prognostic Scores improves the prediction of outcome in Community-Acquired Pneumonia; *Thorax* doi:10.1136/thoraxjnl-2013-203623 <https://www.researchgate.net/>
13. McNally JD, Manaseki-Holland S. Vitamin D deficiency in young children with severe acute lower respiratory infection. *Pediatric Pulmonology*, 2009, 44(10):981–988
14. Quraishi SA, Bittner EA, Christofer KB, Camargo CA Jr. Vitamin D status and Community-Acquired Pneumonia: Results from the Third National Health and Nutrition Examination Survey-NHANES; *PLoS ONE*, 2013; 8(11): e81120. doi: 10.1371/journal.pone.0081120
15. Pletz MW, Terkamp C, Schumacher U, Rohde G, Schütte H, Welte T, Bals R; CAPNETZ-Study Group. Vitamin D deficiency in community-acquired pneumonia: low levels of 1,25(OH)₂ D are associated with disease severity. *Respir Res.* 2014; 15:53. doi: 10.1186/1465-9921-15-53
16. Camargo CA Jr, Ganmaa D, Frazier AL, Kirchberg FF, Stuart JJ, Kleinman K, Sumberzul N, Rich-Edwards JW. Randomized trial of vitamin D supplementation and risk of Acute Respiratory Infection in Mongolia; *Pediatrics* 2012;130-133; e561
17. Bergman P, Lindh ÅU, Björkhem-Bergman L, Lindh JD. Vitamin D and Respiratory Tract Infections: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *PLoS ONE* 8(6): e65835. journals.plos.org/plosone/article?id=10.1371/journal
18. Manaseki-Holland S, Maroof Z, Bruce J, Mughal MZ, Masher MI, Bhutta ZA, Walraven G, Chandramohan D. Effects of vitamin D supplementation to children diagnosed with pneumonia in Kabul: a randomised controlled trial. *Trop Med Int Health* 2010; 15:1148-55
19. Choudhary N, Gupta P. Vitamin D supplementation for severe pneumonia-A randomized controlled trial. *Indian Pediatr.* 2012; 49:449-454.
20. Bryce J, Boschi-Pinto C, Shibuya K, Black RE. WHO estimates of the causes of death in children. *Lancet.* 2005; 365:1147–1152

How to cite this article: Priya PS, Elizabeth KE. Nutritional factors and vitamin d status in relation to severity of pneumonia among children. *Int J Health Sci Res.* 2017; 7(6):6-10.
