

Original Research Article

Somatometric Estimation of Stature Using Head Length and Height Measurements in Male and Female Youths from a Rural Population in Mangalore

Pramod KL¹, Vaswani VR², Khan F³

¹Curator, ²Professor & Head, ³Post Graduate,
Department of Forensic Medicine & Toxicology, Yenepoya Medical College, Mangalore.

Corresponding Author: Khan F

ABSTRACT

Studies done by earlier researchers have shown the importance of stature in the determination of race, age, and sex especially when disaster strikes. In countries like India which shows diversity not only in culture, religion but also in the physical appearance of Individuals which vary from state to state, makes it important to have age and sex specific stature estimation formulae. There is no study based on Mangalore population. The present study was an attempt to find out the stature of an individual from head length (Glabella to Opisthocranium) and head breadth (Euryon to Euryon) measurements and to derive regression formulae to estimate the height in the rural population (age 18-21 years) in Mangalore. It was found that for every unit increase in head length there was a 1.491 unit increase in stature in females and for every unit increase in head breadth there was 0.972 unit decrease in stature. In the males for every unit increase in head length there was a 3.36 unit increase in stature and for every unit increase in head breadth there was 0.962 unit decrease in stature

Key words: Glabella, Opisthocranium, Euryon, Stature

INTRODUCTION

The height of an individual has always been important in the estimation of stature when seen from the eyes of Forensic expert, an Anthropologist or an Archaeologist. ⁽¹⁾ The importance becomes even more evident at the time of Mass disaster's like Earthquake, Flood/Tsunami or air crashes when identifying individuals becomes the most important and difficult task. The knowledge of head dimensions of individuals also helps in determining to which different racial, geographic background they belong to and is specific to that country. ^(2,3) Studies have shown the need for race, age, and sex specific stature estimation formulae. ^(4,5) In India not only the language, but also the physical features of individuals like height, skin color vary as

we move from one state to another, unlike other countries, the diversity seen in Indian population makes it important to have age and sex specific stature estimation formulae. ^(6,7) There is no study based on Mangalore population, This study is a step taken to estimate the stature of an individual from a rural population in Mangalore using the head length and head breadth so as to formulate population and gender specific regression equation.

RESEARCH QUESTION

Can the distance between Glabella and Opisthocranium help in estimating the stature of males and females in a rural population in Mangalore?

OBJECTIVE-

To estimate the stature from head length and head breadth in males and females.

MATERIALS AND METHODS

This study was done as a part of the Post Graduate Diploma in Forensic Anthropology & Odontology (PGDFAO) conducted annually by Forensic Medicine Department of Yenepoya University, Mangalore.

Sample Size-

167 participants (81 males & 86 females) from the age group of 18-21 years using the formula

$$n = \frac{(Z_{1-\alpha/2} + Z_{1-\beta})^2}{[FZ(\rho_1) - FZ(\rho_0)]^2} + 3$$

$$FZ(\rho_1) = 1/2 \ln \left[\frac{1 + \rho_1}{1 - \rho_1} \right]$$

$$FZ(\rho_0) = 1/2 \ln \left[\frac{1 + \rho_0}{1 - \rho_0} \right]$$

ρ_0 = Population Correlation coefficient

ρ_1 = Sample Correlation coefficient

$Z_{1-\alpha/2}$ = Desired confidence level

$1 - \beta$ = Power

Sample correlation Coefficient: 0.93

Population Correlation: 0.9 (Ref 8: Singh R. Estimation of stature and age from head dimensions in Indian population. *Int. J. Morphol.*, 2013;31(4):1185-1190.)

Power : 80%

Level of Significance: 5 % (significance $p < 0.05$)

METHODOLOGY

The study commenced after getting ethical clearance from Yenepoya University Ethics Committee. The participants in the study were 167 Students between the age group of 18 to 21 from two colleges in Mangalore, selected irrespective of their caste and creed. The students were native of Bantwal, Belthangady, Kotekar, Harekala Pavor in Mangalore. The participant information sheet was given to each student

and their role in the study was explained to them. All the participants signed the informed consent form and then the data was collected.

INCLUSION CRITERIA-

All the healthy males and females between the age group of 18 - 21 years from rural population in Mangalore.

EXCLUSION CRITERIA-

People suffering from congenital disorders, dwarfs, giants or having any other physical deformation.

First stature was measured in centimeter unit using the stadiometer (from vertex to floor in Frankfurt's horizontal Plane) using all the standard precautionary measures and the reading was recorded (fig 1).




Fig 1: Measurement of stature using stadiometer. 1- Back of the head, 2- FH plane, 3- Shoulder blade, 4- Bottom, 5- Back of the leg and 6- Heel (<https://www.slideshare.net/reinaramesh/basic-anthropometry-ppt>)

For the measurement of head length (from Glabella to Opisthocranium) and Head breadth (from Euryon to Euryon) spreading caliper (Fig 2, 3 & 4) was used.

Before taking the measurements a straight line was drawn from the trignon to the infra orbital margin using a lip liner on the participant to ensure they are in the Frankfurt's Horizontal Plane. The data collected was then analyzed by mean, standard deviation, Correlation coefficient and by regression analysis.




Fig 2 Head Breadth Measurement (Euryon to Euryon)

Euryon




Fig 3 Head Length Measurement (Glabella to Opisthocranium)

- Glabella
- Opisthocranium



Fig 4- Spreading caliper with rounded ends, Researcher taking Head length measurement

Statistical Analysis:

The data collected was analyzed using IBM SPSS 22. Descriptive statistics was reported as Mean (SD) for continuous variables. Simple linear regression analysis was used to to predict stature from Head length and Head breadth for males and females separately.

RESULT

Total numbers of participants were 167, the distribution of male and female is as shown in Fig 5.




Fig 5: Pie chart showing gender distribution of participants

Table no.1 Descriptive statistics in male with respect to stature, head length & head breadth

Descriptive Statistics					
MALE	n	Minimum	Maximum	Mean	Std. Deviation
Stature (in cm)	81	149.0	178.0	163.253	6.8405
Head Length(in cm)	81	13.0	19.5	16.738	1.0694
Head Breadth (in cm)	81	11.0	16.5	13.094	1.1542
Age	81	18.0	23.0	18.901	.9435

Descriptive statistics for stature, head length, head breadth among males and females is shown in Table 1 and 2

The mean height in both the genders was significantly different and head length, head breadth was found to be larger in males compared to females. Correlation coefficient between the stature, head length and head breadth in both the genders was found to be statistically significant.

In males the stature of the participant was in the range of 149-178, mean height is

163.2 with Standard deviation 6.84. The mean head length and head breadth were 19.5, 16.5 with standard deviation 1.0 & 1.1

Regression equation to predict stature in males

$$\text{Stature} = 119.525 + (- 0.962) \text{ Head breadth} + (3.365) \text{ Head length}$$

For every unit increase in head length there is a 3.36 unit increase in stature and for every unit increase in head breadth there is 0.962 unit decrease in stature

Table no. 2 Descriptive statistics in female with respect to stature, head length & head breadth

Descriptive Statistics					
FEMALE	n	Minimum	Maximum	Mean	Std. Deviation
Stature (in cm)	86	142.5	169.5	154.213	5.4543
Head Length(in cm)	86	13.0	17.5	15.669	.9475
Head Breadth (in cm)	86	10.0	13.5	12.109	.7773
Age	86	17.0	21.0	19.163	.8382


In females the stature of the participant was in the range of 142.5 -169.5. The mean height was 154.2 with a standard deviation of 5.4544. The mean head length and head breadth was 15.6, 12.10 with standard deviation 0.9 & 0.77

Regression equation to predict height in Females

$$\text{Stature} = 142.619 + (- 0.972) \text{ Head breadth} + (1.491) \text{ Head length}$$


For every unit increase in head length there is a 1.491 unit increase in stature in females and for every unit increase in head breadth there is 0.972 unit decrease in stature

SCATTER PLOT SHOWING CORRELATION BETWEEN HEAD LENGTH AND STATURE IN FEMALES



r = 0.199 p = 0.04

SCATTER PLOT SHOWING CORRELATION BETWEEN HEAD LENGTH AND STATURE IN MALES



r = 0.424 ; p ≤ 0.001

Table no. 3 showing comparison between stature, head length and head breadth in Males and females

GENDER	STATURE				HEAD LENGTH				HEAD BREADTH						
	Mean	±	Std	t	p	Mean	±	Std	t	p	Mean	±	Std	t	P
MALES	163.25	±	6.84	9.470	0.000	16.73	±	1.069	6.85	0.000	13.09	±	1.15	6.46	0.000
FEMALES	154.21	±	5.45	9.407	0.000	15.66	±	0.947	6.82	0.000	12.10	±	0.777	6.39	0.000

$P \leq 0.001$, It is significant in both males and females with respect to stature, head length and head Breadth

DISCUSSION

The biological profile of an individual is of great importance in the field of Anthropology and stature estimation is one of the most relevant parameters in building it. (8) Head length measurement by Kadagoudar (2014) was taken from nasion to inion, (9) Jadhav (2004) had measured from glabella to Inion (6) while in the present study the head length was measured from glabella to opisthocranium as it gives the maximum cranial length (ref Martin & Saller, 1957)). There is no standard method to find the relationship between head length and stature as shown by various studies done by researchers in India and abroad, giving a different regression formula for the selected population showing the need for developing population specific regression formula in a country like India. (10-12)

Only other studies to use the head length measurement from glabella to opisthocranium was done by Kumar (2013) and Wankhede (2015). (13,14)

Anthropologist should take an initiative to develop the regression formulae for population in every state of India which can be used for identification in times of disaster. In the study done by Akhtar et al (2009) and Mansur (2014) only head circumference showed significant positive correlation with stature. (1,2) The range of height seen in this study was (149 - 178) cm in males and (142.5 -169.5) cm in females, head length was in the range of (13- 19.5) cm in males and (13 – 17.5) cm in females and head breadth was from (11 – 16.5) cm in males and (10 – 13.5) cm in females.

Table no 4: Comparison of mean height, mean headlength and head breadth with other studies (3,5,8-10,14,15)

Researcher	SS/Age/SP	MHwith SD	MHLwith SD	MHBwith SD	CC (r)
Singh, R (2013)	208 17-26 yrs UttarPradesh	M - 167±7.8 F- 157±0.6	Gla - Ini M- 17±0.7 F- 16±0.8	Lat.pts on P B M- 13.1±0.6 F- 12.5±0.6	M- 0.74 F- 0.0069
Kumar M (2013)	800 >18 yrs Haryana	---	Gla-OpCr M-18.75±1.38 F- 17.75±0.84	Max BPD M-13.11 ±1.09 F- 12.95±0.83	M-0.17 F – 0.19
Chaurasia (2014)	506 18- 25 yrs Indore	M-170.9±11.3 F-153.3±20.5	Gla-Ini M- 18.4±1.2.7 F- 17.26±0.59	---	M – 0.24 F – 0.17
Kadagoudar (2014)	200 19-28 yrs South Indians	M- 172.44 F- 159.49	Nasi- Ini M– 18.18 F- 16.77	---	M – 0.50 F– 0.44
Agarwal (2014)	800 17-25yrs North Indians	M- 169.4±6.04 F- 156.93±5.05	M- 18.24±0.64 F- 17.34±0.62	M-14.86±0.71 F- 14.13±0.54	M - 0.21 F – 0.34
Prasad A 2014	250 18-28 yrs Maharashtra	M-171.93±5.9 F-165.43±3.0	Gla- Ini M-19.76±1.57 F-18.28±1.13	---	M – 0.26 F- 0.22
Wankhede (2015)	470 18-24 yrs Central Indians	M- 170.9±6.8 F- 156.89±5.8	Glab- OpCr M-18.5±0.72 F- 17.53±0.81	Eu– Eu M-14.64±0.63 F- 14.16±0.66	M – 0.27 F- 0.20
Present study (2015)	167 18 – 21 yrs Mangalore	M– 163.2±6.8 F - 154±5.4	Glab- OpCr M- 16.7±1.06 F- 15.6±0.94	Eu – Eu M- 13.09±1.15 F- 12.1±0.77	M – 0.42 F – 0.19

(SS- Sample size, SP- Study Population, MH – Mean Height, SD- Standardd Deviation, MHL- Mean head length, MHB- Mean head breadth, CC- Correlation coefficient, Glabella-Inion- GI, Lat.pts on P B – Lateral points on parietal bones, Maximum Biparietal diameter- MBPD, Euryon – Euryon- Eu- Eu, Glabella –Opisthocranium – Gla- Op Cr, Nasion – Inion Nasi-Ini, M- Male, F- Female).

The mean stature in male was 163.25 ± 6.84 and females was 154.21 ± 5.45 , mean head length in males was 16.73 ± 1.069 while in females it was 15.66 ± 0.947 and mean head breadth in males was 13.09 ± 1.15 which was more compared to the females mean head breadth 12.10 ± 0.777 . The p value was significant for both the genders with respect to stature, head length and head breadth, which is same as a study of North Indian Populations by Agarwal (2014) showing correlation coefficient between stature and head length and head breadth significant (10). In the present study the correlation between stature and head length in males was $r = 0.424$, $p \leq 0.001$ in males and $r = 0.199$, $p \leq 0.001$ in females when compared to studies done by Mansur (2014) which showed the value of $r = 0.0443$, $p < 0.001$ in males and $r = 0.302$, $p < 0.001$ in females (Nepalese population), (2) by Chaurasia (2014) in Indore correlation coefficient was 0.241 in males and 0.173 in females, (3) a study done by Kumar (2013) in Rajasthan population showed correlation coefficient was 0.941 in males and 0.85 in females. (5) The correlation coefficient in Gujarat population in a study done by Jadhav (2004) was 0.53, (6) Kadagoudar (2014) studied South Indian population and found $r = 0.507$ in males and 0.440 in females, (9) while in a study done by Agarwal (2014) $r = 0.215$, $p < 0.001$ in males and $r = 0.341$, $p < 0.001$ in females in North Indian Population. (10)

The correlation coefficient of Stature and head length of females (0.19) in the present study is same as the study done by Kumar (2013) in Haryanavi's (13) while it is slightly lower than the study done by Wankhede (2015) which was 0.20. (14) In males the correlation coefficient 0.42 of the present study was lower than the study done by Kadagoudar (2014) which was 0.50 (9) but in the present study head length was measured from glabella to opisthocranium while in the latter it was taken from nasion to inion.

There is a need for conducting research to find gender specific regression

equations for the various populations in different states of India. The measurements for stature, head length, head breadth should be taken from standardized landmarks as done by anthropologists. The researcher should be either an anthropologist or one with training in anthropology to do justice to the study.

SUMMARY & CONCLUSION

- Head length measured from Glabella to Opisthocranium showed positive correlation with that of stature in both Males and females.
- The regression equation for Males is $119.525 + (-0.962) \text{ Head breadth} + (3.365) \text{ Head length}$, For every unit increase in head length there is a 3.36 unit increase in stature and for every unit increase in head breadth there is 0.962 unit decrease in stature
- The regression equation for Females is $142.619 + (-0.972) \text{ Head breadth} + (1.491) \text{ Head length}$, For every unit increase in head length there is 1.491 unit increase in stature in females and for every unit increase in head breadth there is 0.972 unit decrease in stature.

Limitations

1. Intraobserver error while taking the head length and breadth
2. Reliability of Instruments used for measurement

Suggestions

1. Standard method of measurement should be used by researchers all over the country. For example – Head length Measurement - Nasion to Inion or Glabella to Inion or Glabella to Opisthocranium (present study).
2. The regression equation which is gender, age, population specific should be derived for the diverse population of India.

ACKNOWLEDGEMENT

I am grateful to the Principals of the two colleges in Mangalore for giving me permission to conduct this study and to all the wonderful, enthusiastic students for their cooperation and support. I would like to extend my gratitude to

Dr. Mohammad Nasir Ahemad, Anthropologist for his valuable suggestions.

I am also thankful to Ms. Soumya Shetty, Biostatistician for helping me with the analysis of the data.

Conflict of Interest: None

Sponsors: None

REFERENCES

1. Akhter Z et al. Stature Estimation Using Head Measurements in Bangladeshi Garo Adult Females. Bangladesh Journal of Anatomy 2009;7(2):101-104.
2. Mansur DI, Haque MK, Sharma K, Mehta DK, Shakya R. Use of Head Circumference as a Predictor of Height of Individual. Kathmandu Univ Med J 2014; 46(2):89-92.
3. Chaurasia RS, Bamne A, Shukla SN, Gohiya V, Pandey R. Estimation of Stature from head length. Inter J Medical Sci Res Prac 2014; 1 (2): 40-43.
4. Shende MR, Tirpude BH, Bokariya P, Kothari R. Significance of Head Length in Estimation of Stature in Healthy Individuals of Central India. 2013. Indian Journal of Forensic Medicine & Toxicology. 2013; 7(2): 128.
5. Kumar S, Garg R, Mogra K, Choudhary R. Prediction of stature by the measurement of head length in population of Rajasthan. Journal of evolution of Medical and Dental sciences 2013;2(14):2334-2340.
6. Jadav HR, Shah GV. Determination of personal height from the length of head in Gujarat Region. J.Anat.Soc.India. 2004; 53 (1):20-21.
7. Umesh SR, Imran SS, Uzair SH. Estimation of stature from headlength in Hyderabad Karnataka region. Medico Legal Update. 2013;13(1):187-189.
8. Singh, R. Estimation of stature and age from head dimensions in Indian population. Int. J. Morphol. 2013; 31(4): 1185-1190.
9. Kadagoudar S, Manjulabai KH, Hallikeri VR. Estimation of stature by Nasion- Inion length in South Indian Population – A cross sectional study. Medica Innovatica 2014 Dec; 3(2):9-11.
10. Agarwal S, Agarwal SK, Jain SK. Correlation between the stature and cranial measurements in population of North India. Acta Medica International 2014 Dec; 1(2):99-102.
11. Ewanonu EO, Anibeze CIP. Estimation of stature from Cephalic parameter in South Eastern Nigerian population. JSIR 2013Apr;2(2):425-432.
12. Bansal H, Badiye A. An estimation of correlation between the Headlength and the stature of the children aged between 6-10 years. Res.J. Forensic Sci. 2013 Apr;1(2):1-5.
13. Kumar M, Gopichand PVV. Estimation of stature from cephalo-facial anthropometry in 800 Haryanvi adults. IJPAES 2013; 3(2):43-46.
14. Wankhede KP, Anjankar VP, Parchand MP, Kamdi NY, Patil ST. Estimation of stature from head length & head breadth in central Indian population: An Anthropometric study. Int J Anat Res 2015;3(1):954-957.
15. Prasad A, Bhagwat VB, Kumar A, Joshi DS. Estimation of stature from head length in living adults in Marathwada region of Maharashtra. Int.J.RSR. 2014; 5(10): 1745-1748.

How to cite this article: Pramod KL, Vaswani VR, Khan F. Somatometric estimation of stature using head length and height measurements in male and female youths from a rural population in Mangalore. Int J Health Sci Res. 2017; 7(8):54-60.
