

Original Research Article

Prevalence of Anemia among Pregnant Women Attending Antenatal Clinic of a Selected Hospital in Accra, Ghana

Kwabena Acheampong¹, Stella Appiah², Dorothy Baffour-Awuah², Yeboah Saka Arhin¹

¹Valley View University Hospital, Public Health and Laboratory Department, P.O.Box AF 595, Adenta-Accra, Ghana

²Valley View University, Faculty of Health Sciences, Department of Nursing, P.O.Box AF 595, Adenta-Accra, Ghana

Corresponding Author: Kwabena Acheampong

ABSTRACT

Background: Anemia troubles about half of Ghanaian grown-up population and its highest value over the past 16 years was 56.40% in 2011, while its lowest value was 49.20% in 1999.

Objective: To determine the prevalence of anaemia among pregnant women attending antenatal clinic of a selected private hospital in Accra, Ghana.

Methodology: Using across-sectional design, 200 consenting participants were selected for the study. Hemoglobin levels were assessed. Maternal age, gestational age, parity and number of antenatal visits were collected from the antenatal record booklet. Data were analyzed using the SPSS programme IBM version 20. Chi-square was used to investigate the association between independent variables and prevalence of anemia. The data were further analyzed using binary logistic regression methods to evaluate possible risk factors associated with anemia. Odd ratio was reported to establish the risk of anemia and 95% confidence interval were estimated. A $p < 0.05$ was considered as statistically significance.

Results: The findings revealed that, the mean with \pm standard deviation of hemoglobin value was 10.9 ± 1.3 (95% [confidence interval] = 10.7–11.1), and the overall prevalence of anemia from a total of 200 study subjects was 102 (51.0%). The severities of all diagnosed anemia cases were mild (60.8%) to moderate (39.2%). Severe anemia was not found in any of the participants.

Conclusion: The study established that the prevalence of anemia was high and it is associated with parity. Preventive efforts targeting preconception nutrition and promotion of regular antenatal care visit throughout the normal pregnancy is urgently required.

Key words: Pregnant Women, Hemoglobin, Trimester, Antenatal Care, Anemia

INTRODUCTION

Globally, the mean blood hemoglobin concentration was 11.1 g/dl. [1] (Stevens et al., 2013). Anemia resulting from iron deficiency adversely affects cognitive and motor development, causes fatigue and low productivity. [2,3] and, when it occurs in pregnancy, may be associated with low birth weight and increased risk of preterm delivery. [4,5,6,7] The mechanisms causal to these effects are unknown, but they may be

related to reduce oxygen delivery to the placenta and fetus, increased rates of infection, or adverse effects of iron deficiency on brain development. [8,9,10]

Anemia is one of the most prevalent nutritional deficiency problems affecting pregnant women. The prevalence of anemia in pregnancy differs significantly because of variations in socioeconomic conditions, lifestyles, and health-seeking behaviors across different cultures. [11,12,13] The World

Health Organization (WHO) estimates that 52% of pregnant women in developing countries are anemic compared with 23% in the developed world. [14] Anemia has also been recognized as one of the greater risk of perinatal mortality and morbidity. [15] Accumulating information suggests an association between maternal iron status in pregnancy and the iron status of infants postpartum. [16,17]

Anemia troubles about half of Ghanaian grown-up population. Its highest value over the past 16 years was 56.40% in 2011, while its lowest value was 49.20% in 1999. [1] The raise in its prevalence will occur largely in pregnant women. Anemia is defined as decrease in the total amount of red blood cells. [18,19] ORA decrease in whole-blood hemoglobin concentration of more than two standard deviations below the mean of an age- and sex-matched reference range. [20] Inadequate folate during pregnancy has long been associated with reduced fetal growth and anemia in pregnancy. [21,22]

Over the years, the proportions of iron-deficiency anemia in pregnancy have lingered high in women in developing as well as developed countries. [23,24,25] Its often occurs toward the end of pregnancy even among women who enter pregnancy with some iron stores. [26,27] It is far more common than iron-deficiency anemia. Iron-deficiency and iron-deficiency anemia are related to reduce iron stores in newborns. [28] The nine months of pregnancy characterize the most intense period of growth and development humans ever experience. How well these processes go depends on many factors, most of which are modifiable. Of the factors affecting fetal growth and development that are within our control to change, nutritional status stands out. [29]

At no other time in life are the benefits of optimal nutritional status more obvious than during pregnancy. Many aspects of nutritional status, such as dietary intake, supplement use, and weight change, influence the course and outcome of pregnancy. [8,9,30] The fetus is not a parasite;

it depends on the mother's nutrient intake to meet its nutritional needs. [31] Periods of rapid growth and development of fetal organs and tissues occur during specific times throughout pregnancy. Vital nutrients must be accessible in required amounts during these times for fetal growth and development to proceed optimally. [32,33]

Pregnancy is a molecule-building process and a woman's normal nutritional requirement increases during pregnancy to meet the needs of the growing fetus and the maternal tissues associated with pregnancy. [34] If energy and other nutrient intake do not enhance, the body's own reserves are used, leaving a pregnant woman weakened. Energy needs increase in the second and particularly the third trimester of pregnancy, mainly due to increased maternal body mass. [35]

The study has the following objectives

1. To determine the prevalence of anemia among pregnant women attending antenatal care in a selected private hospital in Accra, Ghana.
2. To determine prevalence of anemia and its association with maternal age, gestational age, parity and number of antenatal care visit.

METHODOLOGY

Study Designs

A cross sectional study included pregnant women aged between 15-44 years attending antenatal clinic in a selected private hospital in Accra, Ghana were eligible to participate in the study. Subjects were informed and their consent was sought. Participants were recruited when they arrived for routine antenatal clinic. Required information on age, level of education, gestational age of the pregnancy, parity and the number of antenatal visits was obtained from interview and antenatal record booklet to gather with current hemoglobin estimation test.

Sample Size and Sampling Method

The sample size was calculated using (Fisher, 1978):

$N = Z^2 * P (1 - P) / d^2$, where N = sample size

Z = score for 95% Confidence Interval which is 1.96

P=prevalence of hypertension (proportion of people with hypertension presume to be 14% or higher).

q=1-p (proportion of people without hypertension) d = tolerable error set at 5%. N =185.01 making 8% allowances for losses, it was approximated to 200.

Sampling Method

Facility based quantitative cross-sectional study was employed from January to August, 2017 at selected private hospital located on Adenta-Dodowa road. The study was designed to be performed over 24 weeks during which estimated 400 women were expected to attend outpatient clinic from January 2016 and August, 2017. The systematic random sampling method was used to select the study participants, at fixed intervals (k = sampling interval: 400/200= 2). Pregnant women who were not lately been given blood or transfused, who had no chronic medical diseases and who had no early bleeding were included in the study. Each participant was recruited only once on their first visit during the study period.

Statistical Analysis

Data were analyzed using the SPSS programme IBM version 20. Descriptive statistics were calculated for all numerical parameter. Continuous variables are stated in mean standard deviation. Cross tabulation were done to determine percentages. Chi-square were used to investigate the association between independent variables and prevalence of anemia. The data were further analyzed using binary logistic regression methods to evaluate variables associated with anemia. Odd ratio (OR) was reported to establish the risk of anemia and 95% confidence interval (C.I.) were estimated. A p<0.05 was considered as statistically significance.

RESULT

Table 1 presents results on the percentage distribution of anemia and

maternal age, level of education, gestational age, parity and number of visit among the study participants Results indicate that moderate anemia was higher among age (35-44)years representing 23.2%. Prevalence of moderate anemia was low among those tertiary (4.8%) as compared to its counterparts. Prevalence moderate anemia was also found to decrease as the gestational age increases, showing the highest prevalence in the first trimester 21.4% than second 20.0% and third trimester 19.8%. Pregnant women with more than four children had shown more prevalence of moderate anemia (23.1%) than its counterparts. Prevalence of moderate anemia was also found to decrease as the participant number of visits increases, showing the lowest prevalence in more than eight visits 11.5%, four to seven visits 20.7% and less or equal to three visit 21.7%.

Table 1 showing the percentage distribution of anemia and maternal age, level of education, gestational age, parity and number of visit among the study participants

Variables	Anemia			No. of subject
	Moderate	Mild	Non	
Maternal Age	Chi-Square =10.737, P=0.097			
15-24	3(15.0)	7(35.0)	10(50.0)	20 (10.0)
25-34	24(19.4)	39(31.5)	61(49.2)	124(62.0)
35-44	13(23.2)	16(28.6)	27(48.2)	56(28.0)
Level of Education	Chi-Square= 0.196, P=0.0978			
Primary & Below	19(20.4)	25(26.9)	49(52.7)	93(46.5)
JHS	15(24.6)	20(32.8)	26(42.6)	61(30.5)
SHS	5(20.0)	12(48.0)	8(32.0)	25(12.5)
Tertiary	1(4.8)	5(23.8)	15(71.4)	21(10.5)
Gestational Age	Chi-Square = 0.703, P= 0.951			
1 st Trimester	3(21.4)	5(35.7)	6(42.9)	14(7.0)
2 nd Trimester	17(20.0)	24(28.2)	44(51.8)	85(42.5)
3 rd Trimester	20(19.8)	33(32.7)	48(47.5)	101(50.5)
Parity	Chi-Square =11.753, P<0.019			
No Child	9(15.5)	11(19.0)	38(65.5)	58(29.0)
1 Child	10(18.9)	16(30.2)	27(50.9)	53(26.5)
2 Children	15(26.8)	20(35.7)	21(37.5)	56(28.0)
3 Children	3(15.0)	9(45.0)	8(40.0)	20(10.0)
≥4.0 Children	3(23.1)	6(46.2)	4(30.8)	13(6.5)
Number of Antenatal Care Visit	Chi-Square =7.014, P=0.135			
≤ 3 times	20(21.7)	30(32.6)	42(45.7)	92(46)
4-7 times	17(20.7)	28(34.1)	37(45.1)	82(41.0)
≥8 times	3(11.5)	4(15.4)	19(73.1)	26(13.0)
1 st , 2 nd & 3 rd Trimester 9(0-12), (13-27) & (28-40) weeks respectively				

Table 2 presents results on the percentage distribution of hemoglobin level among the study participants. In terms of gravity, mild anemia was 60.8%, moderate anemia was 39.2% and severe anemia was 0.0%.

Table2: Percentage distribution of Hemoglobin level among the study population.

Hemoglobin level(g/dl)	Percentage	No. of Subject
Severe Anemia (≤ 7)	0.0%	0
Moderate Anemia (7-9.9)	39.2%	40
Mild Anemia(10.10.9)	60.8%	62
Non-Anemia (≥ 11)	49.0%	98

Table3: Mean Hemoglobin estimation among Pregnant Women

Gestational Age	Mean \pm Std. Deviation	No. of Subject
1 st Trimester	10.643 \pm 1.2283	14
2 nd Trimester	10.959 \pm 1.3924	85
3 rd Trimester	10.895 \pm 1.2956	101

Prevalence of Anemia

The mean \pm SD hemoglobin concentration was 10.9g/dl \pm 1.33 and an overall prevalence rate of anemia with hemoglobin level <11 g/dl was 51.0% (CI: 10.7-11.1).

Table 4: Biological data and Interpreting Hemoglobin

Variable	Range	Mean	Std. Dev
Hemoglobin ((g/dl))	7.3 - 15.3	10.90	1.33
Interpreting hemoglobin	Anemia	Number	%
	< 11	102	51.0%
	≥ 11	98	49.0%
			Standards
			12-16

A person with hemoglobin below <11 g / dl is anaemic according to World Health Organisation. [1]

Binary Logistic Regression Risk Factors Analysis for Anemia

The results of binary logistic regression analysis, including the odds ratio for maternal age, gestational age, parity and number of visit are presented in table 8. Significant association was related to 4-7 times antenatal visit ($p < 0.006$), other factors like maternal age, gestational age, parity were also analyzed but no significant associations were found.

Inferential Statistics

Table 5 shows that participants who were aged 35-44 were 1.057 times more likely to be anemic than those who were below the age of 35 years. However the rate of moderate anemia increases with increases in age and vice versa from mild anemia as shown in table 1. The gestational age between 28-40 weeks were 1.832 times more likely to be anaemic than those who

were below. Participants with three or more children were 1.874 times more likely be anaemic than those who were below. The number of antenatal visits revealed no positive association with prevalence of anemia.

Table 5: Association between prevalence anemia and maternal age, gestational age, parity and number of visit

Parameters	Odd ratio [95% CI]	p-value
Age (years)		
15-24	1.0	
25-34	0.971 [0.361-2.610]	0.954
35-44	1.057 [0.357-3.128]	0.920
Gestational Age		
0-12	1.0	0.258
13-27	1.395 [0.383-5.082]	0.614
28-40	1.832 [0.884-3.797]	0.104
Parity		
≤ 3 Children	1.0	
≥ 3 Children	1.874[0.823-4.267]	0.135
No. of Visit		
0-3	1.0	
4-7	0.189[0.061-0.592]	0.006
≥ 8	0.244[0.089-0.667]	0.135

DISCUSSION

Prevalence of Anemia

Antenatal care allows management of pregnancy, detection and treatment of complications, and promotion of better maternal and child health. [36] However, women rarely recognize childbearing as problematic and, therefore; do not seek care. [37] This study recognized anemia as a major health problem among pregnant women in Ghana. In the *present study*, 51% of the participants are found to be anemic.

This figure is slightly lower compared to World Bank report in 2016 on prevalence of anemia which was given as 54% among pregnant women in Ghanaian population. [1] Similar results were reported on prevalence of anemia among pregnant women in previous studies. [38,39] The higher prevalence may be due to inadequate dietary iron intake and lack of proper nutrition during preconception period. In this study, about one in every two pregnant women experiences iron-deficiency anemia during pregnancy. The present study indicated that moderate anemia (39.2%) constitutes a significant portion of anemia in pregnant women attending the antenatal clinics.

Maternal Age and Anemia

Statistical significance association between parity and anemia was found ($p < 0.097$). This is inconsistent with Chowdhury and colleagues who reported association of anemia with maternal age. [13] When maternal age was considered (Table 1), we observed that moderate anemia increases along with increases maternal age and vice versa for mild anemia.

Gestational Age and Anemia

Gestational age of the pregnancy were important parameters but have revealed no statistical significance association with anemia in the current study ($P = 0.951$). Some studies have reported results of the association between gestational age and anemia. [40,41,42] The present study revealed high proportions of mild and moderate anemia combine in first trimester and reaches the lowest rate at the end of second trimester and increase in third trimesters as indicated in table 1.

This finding is consistent with a study conducted by Vemulapali and Roa, 2014 in Andhra Pradesh reported that pregnant women in first trimester showed higher anemia prevalence rate than in second and third trimesters. [43] But inconsistent with a study which reported on prevalence of anemia among pregnant women in rural population of Kolar in district. [44]

As pregnancy progresses, plasma volume progressively increases and most of this increase occurs by 34 weeks' gestation and is proportional to the birth weight of the baby. Because the expansion in plasma volume is greater than the increase in red blood cell mass, there is a fall in haemoglobin concentration, haematocrit and red blood cell count. [28,45]

Parity and Anemia

Statistical significance association between parity and anemia was found ($p < 0.019$). Similar result was reported on association of anemia with parity. [46] Concerning parity (Table 1), it was observed in this study that moderate anemia was less common (15.5%) in primigravidas as

compare to 18.9%-26.8% for multiparas, while mild anemia followed the same trend, 19.0% anemia in primigravidas and 30.2%-46.2% anemia in multiparas.

Several studies have observed higher prevalence of anemia in women with high parity. [14,46] Likely rationalization to the high prevalence of anemia among multiparous women is that these women might have gotten pregnant with low levels of nutrients due to the reduction of reserves of the mother in prior pregnancies and lactation periods or possibly did not also allow spaces between their pregnancies so were not fully recovered from previous anemia.

Number of Antenatal Care Visit and Anemia

Statistical significance association between number of antenatal visit and anemia was not found ($P = 0.135$). Regarding number of antenatal visit (Table 1) it was observed in this study that prevalence of moderate anemia decreased with increased number of visit. This indicates that regularly routine checked of hemoglobin might have prompted the women to eat well and also supplements that are given monthly might have played significant role in reducing anemia in pregnant women who visit regularly.

The results of our study showed that level of education of mothers had a significant association with maternal anaemia. This finding is supported by another study in which literacy of women had a significant association with the use of antenatal care services, as education has an impact on awareness of use of health services among the population. [47]

Mean Hemoglobin estimation among Pregnant Women

The current study demonstrates that mean hemoglobin levels decreased during the first trimester and attained its highest peak at the end of second trimester, and decreases again during the third trimester of pregnancy as shown in table 3. This finding is inconsistent with a research that has revealed that hemoglobin concentrations

typically decrease during the first trimester and reach the lowest levels at the end of second trimester and increase again during the third trimester of pregnancy. [48] It is evident that, the significant declined of hemoglobin level in pregnant women is due in part to dietary iron deficiency, therefore iron therapy is helpful to maintain hemoglobin nearer to that of normal women. [49]

CONCLUSION

- The study established high prevalence of anemia and it was associated with parity
- Hemoglobin appears to increase with advancing pregnancy, which suggests the mild risk of anemia in pregnancy
- Prevalence of anemia decrease with increase number of antenatal care visit.

RECOMMENDATION

Pregnant women should be educated on adequate healthy dietary intake to maintained normal hemoglobin levels. In a unique effort to prevent this problem, pregnant women should encourage to attending antenatal care more often to decrease the incidence of anemia. Finally women should be advised to take 30–60 mg of iron along with folic acid in a supplement.

REFERENCES

1. Stevens GA, Finucane MM, De-Regil LM, Paciorek CJ, Flaxman SR, Branca F et al. Global, regional, and national trends in haemoglobin concentration and prevalence of total and severe anaemia in children and pregnant and non-pregnant women for 1995–2011: a systematic analysis of population-representative data. *Lancet Glob Health*. 2013;1:E16–E25. doi:10.1016/S2214-109X(13)70001-9.
2. Balarajan Y, Ramakrishnan U, Ozaltin E, Shankar AH, Subramanian SV. Anaemia in low-income and middle-income countries. *Lancet*. 2011;378:2123–35. doi:10.1016/S0140-6736(10)62304-5.
3. Veena SR, Gale CR, Krishnaveni GV, Kehoe SH, SrinivasanK, & Fall CH. Association between maternal nutritional status in pregnancy and offspring cognitive function during childhood and adolescence; a systematic review. *BMC Pregnancy and Childbirth*. 2016; 16(1), 220.
4. Brabin L, Brabin BJ, & Gies S. Influence of iron status on risk of maternal or neonatal infection and on neonatal mortality with an emphasis on developing countries. *Nutrition Reviews*.2013;71(8), 528-540.
5. Kozuki N, Lee AC, Katz J, Child Health Epidemiology Reference Group. Moderate to severe, but not mild, maternal anemia is associated with increased risk of small-for-gestational-age outcomes. *J Nutr*. 2012;142:358–62. doi:10.3945/jn.111.149237.
6. Rao R, & Georgieff MK. Iron in fetal and neonatal nutrition. In *Seminars In Fetal And Neonatal Medicine*. 2007;12(1):54-63. WB Saunders.
7. Kavle JA, Stoltzfus, RJ, Witter F, Tielsch JM, Khalfan SS, & Caulfield, LE. Association between anaemia during pregnancy and blood loss at and after delivery among women with vaginal births in Pemba Island, Zanzibar, Tanzania. *Journal of Health, Population, and Nutrition*.2008; 26(2), 232.
8. Grieger JA, & Clifton VL. A review of the impact of dietary intakes in human pregnancy on infant birth weight. *Nutrients*, 2014; 7(1), 153-178.
9. Darnton-Hill I, & Mkparu UC. Micronutrients in pregnancy in low-and middle-income countries. *Nutrients*.2015; 7(3), 1744-1768.
10. Leventakou V, Roumeliotaki T, Martinez D, Barros H, Brantsaeter AL, Casas M, ... &Forastiere F. Melku M, Addis Z, Alem M, Enawgaw B. Prevalence and Predictors of Maternal Anemia during Pregnancy in Gondar, Northwest Ethiopia: An Institutional Based Cross-Sectional Study. *Anemia*. 2014; 2014:108593. doi:10.1155/2014/108593.
11. MulugetaMelku, Zelalem Addis, Meseret Alem, and Bamlaku Enawgaw, “Prevalence and Predictors of Maternal Anemia during Pregnancy in Gondar, Northwest Ethiopia: An Institutional Based Cross-Sectional Study,” *Anemia*, vol. 2014, Article ID 108593, 9 pages, 2014. doi:10.1155/2014/108593Singh P; Khan & S; Mittal RK. anemia during pregnancy in the women of western Nepal. *Bali Medical Journal*, [S.l.], Jan. 2013. ISSN 2302-2914. Available at: <<https://ojs.unud.ac.id/index.php/bmj/article/view/5250>>.
12. Zehra T, Khan RA and, QadriF .*Anemia in Pregnancy a Study at Karachi in a Tertiary. Care Centre*. American Journal of Phytomedicine and Clinical Therapeutics, 2017; 1224-1233
13. Chowdhury HA, Ahmed KR, Jebunessa F, Akter J, Hossain S, Shahjahan M. Factors

- associated with maternal anaemia among pregnant women in Dhaka city. *BMC Women's Health*. 2015;15:77. doi:10.1186/s12905-015-0234-x.
14. Al-Mehaisen L, Khader Y, Al-Kuran O, Abu Issa F, & Amarín, Z. (Maternal Anemia in Rural Jordan: Room for Improvement. *Anemia*, 2011, 381812. <http://doi.org/10.1155/2011/381812>
 15. Abu-Ouf NM, Jan MM. The impact of maternal iron deficiency and iron deficiency anemia on child's health. *Saudi Medical Journal*. 2015;36(2):146-149. doi:10.15537/smj.2015.2.10289.
 16. Schantz-Dunn J, Nour NM. Malaria and Pregnancy: A Global Health Perspective. *Reviews in Obstetrics and Gynecology*. 2009;2(3):186-192.
 17. Khan S. A study of maternal and foetal outcome In maternal anemia.2012. Navali N. Assessment of relationship between Iron deficiency and preterm labor. *Life Science Journal*, 2012; 9(4).
 18. Olatunbosun OA, Abasiattai AM, Bassey EA, James, RS, Ibanga, G, & Morgan A. Prevalence of anaemia among pregnant women at booking in the University of Uyo Teaching Hospital, Uyo, Nigeria. *BioMed Research International*, 2014.
 19. Maakaron, Joseph "Anemia: Practice Essentials, Pathophysiology, Etiology". *Emedicine*. Archived from the original on 14 October 2016. Retrieved C0 October 2017.
 20. Pomeranz, AJ, *Pediatric Decision-Making Strategies (2nd ed.)*, Elsevier,(2016), ISBN 978-0323298544.
 21. Greenberg JA, Bell SJ, Guan Y, & Yu YH. Folic acid supplementation and pregnancy: more than just neural tube defect prevention. *Reviews in Obstetrics and Gynecology*, 2011; 4(2), 52.
 22. Bodnar LM, Himes KP, Venkataramanan R, Chen JY, Evans RW, Meyer JL, & Simhan HN. Maternal serum folate species in early pregnancy and risk of preterm birth. *The American Journal of Clinical Nutrition*.2010; 92(4), 864-871.
 23. Baig-Ansari N, Badruddin SH, Karmaliani R, Harris H, Jehan I, Pasha O,..& Goldenberg RL. Anemia prevalence and risk factors in pregnant women in an urban area of Pakistan. *Food and Nutrition Bulletin*, 2008; 29(2), 132-139.
 24. Alem M, Enawgaw B, Gelaw A, Kena T, Seid M, & Olkeba Y. Prevalence of anemia and associated risk factors among pregnant women attending antenatal care in Azezo Health Center Gondar town, Northwest Ethiopia. *Journal Interdiscipl Histopathol*. 2013; 1(3), 137-144.
 25. Kefyalew Addis Alene and Abdulahi Mohamed Dohe. Prevalence of Anemia and Associated Factors among Pregnant Women in an Urban Area of Eastern Ethiopia. *Anemia*. vol. 2014, Article ID 561567, 7 pages, 2014. doi:10.1155/2014/561567
 26. Kalaivani K. Prevalence and consequences of anemia in pregnancy. *Indian J Med Res*.2000; 130: 627-33
 27. McMahon LP. Iron deficiency in pregnancy. *Obstetric Medicine*. 2010;3(1), 17-24.
 28. Terefe B, Birhanu A., Nigussie P., & Tsegaye A. Effect of maternal iron deficiency anemia on the iron store of newborns in Ethiopia. *Anemia*, 2015.
 29. Judith E. Brown, Janet S. Isaacs, U. Beate Krinke, Ellen Lechtenberg, Maureen A. Murtaugh, Carolyn Sharbaugh, Patricia L. Splett, Jamie Stang, Nancy H. Nutrition Through the Life Cycle 4th ed 3rd ed Wooldridge(2011). *Wadsworth. Cengage Learning*.
 30. Colón-Ramos U, Racette SB, Ganiban J, Nguyen TG, Kocak M, Carroll KN., ... & Tyllavsky FA. Association between dietary patterns during pregnancy and birth size measures in a diverse population in Southern US. *Nutrients*, 2015;7(2), 1318-1332.
 31. Marangoni F, Cetin I, Verduci E, Canzone G, Giovannini M, Scollo P, ... & Poli A. Maternal Diet and Nutrient Requirements in Pregnancy and Breastfeeding. An Italian Consensus Document. *Nutrients*, 2016;8(10), 629.
 32. Hay WW. Aggressive Nutrition of the Preterm Infant. *Current pediatrics reports*. 2013;1(4):10.1007/s40124-013-0026-4. doi:10.1007/s40124-013-0026-4.
 33. Greer FR, & Olsen IE. (2013). How fast should the preterm infant grow?. *Current Pediatrics Reports*. 2013; 1(4), 240-246.
 34. Durrani AM, & Rani A. Effect of maternal dietary intake on the weight of the newborn in Aligarh city, India. *Nigerian Medical Journal: Journal of the Nigeria Medical Association*, 2011;52(3), 17
 35. Diemert A, Lezius S., Pagenkemper M, Hansen G, Drozdowska A, Hecher K.,... & Zyriax BC. Maternal nutrition, inadequate gestational weight gain and birth weight: results from a prospective birth cohort. *BMC Pregnancy and Childbirth*, 2016; 16(1), 224.
 36. Al-Ateeq MA, Al-Rusaies AA. Health education during antenatal care: the need for more. *International Journal of Women's*

- Health.2015;7:239-242
doi:10.2147/IJWH.S75164.
37. Md. Shahjahan, Hasina Akhter Chowdhury, Jesmin Akter, AfsanaAfroz, M Mizanur Rahman, MA Hafez. Factors associated with use of antenatal care services in a rural area of Bangladesh .South East Asia Journal of Public Health 2012;2(2):61-66.
 38. Jagadish Kumar K, Asha N, Srinivasa Murthy D, Sujatha MS, and Manjunath VG. Maternal anemia in various trimesters and its effect on newborn weight and maturity: An observational study.Int J Prev Med. 2013;4(2): 193-199
 39. Jufar AH, Zewde T. Prevalence of Anemia among Pregnant Women Attending Antenatal Care at Tikur Anbessa Specialized Hospital, Addis Ababa Ethiopia. J Hematol Thromb Dis.2014; 2:125. doi:10.4172/2329-8790.1000125
 40. Bencaiova G, & Breymann C. Mild anemia and pregnancy outcome in a Swiss collective. Journal of Pregnancy, 2014.
 41. Yildiz Y, Özgü E, Unlu SB, Salman B, &Eyi EGY. The relationship between third trimester maternal hemoglobin and birth weight/length; results from the tertiary center in Turkey. The Journal of Maternal-Fetal & Neonatal Medicine.2014 27(7), 729-732.
 42. Ren A, Wang J, Ye RW, Li S, Liu JM, & Li Z. Low first-trimester hemoglobin and low birth weight, preterm birth and small for gestational age newborns. International Journal of Gynecology & Obstetrics. 2007; 98(2), 124-128.
 43. Ibhargavi Vemulapalli & Kameswara Rao. Prevalence of anemia among pregnant women of rural community in Vizianagram, North Coastal Andhra Pradesh, India. Asian Journal of Medical Science, 2014 Volum5 <http://nepjol.info/index.php/AJMS>
 44. Suryanarayana R, Santhuram, AN, Muninarayana Chandrappa, PS, & Rangappa SS. Prevalence of anemia among pregnant women in rural population of Kolar district. International Journal of Medical Science and Public Health. 2016; 5(3), 454-458.
 45. Bassi Amos Paul, Idoko Lucy, Dibigbo-IbeajiNdudi Mary, AdeniyiOlusegun Godwin, RamyilMamzhi-Crown Seljul, Ogundeko Timothy Olugbenga, Pisagih James Kurgnan, Offiah Valeria Nneka, Olorunfunmi Joshua Seye, Amuta Wilson Ehi, Alu Vivian Ojoma, AjalaEse Kate, Olorundare Olufunke Olufunmilola. Prevalence of Anaemia in Pregnancy Among Women Visiting Antenatal Clinic in Bingham University Teaching Hospital Jos, Nigeria. Clinical Medicine Research. 2016;5(3):52-62. doi: 10.11648/j.cmr.20160503.17
 46. Taner CE,Ekin A, Solmaz U, Gezer C, Çetin B, KeleşoğluM, ...Özeren M. (2015). Prevalence and risk factors of anemia among pregnant women attending a high-volume tertiary care center for delivery. Journal of the Turkish German Gynecological Association.2015;16(4), 231-236. <http://doi.org/10.5152/jtgga.2015.15071>
 47. Yahya M Al-Farsi, Daniel R Brooks Martha M Werler, Howard J Cabral, Mohammed A Al-Shafei† and Henk C Wallenburg. Effect of high parity on occurrence of anemia in pregnancy: a cohort study. BMC Pregnancy and Childbirth. 2011; 11:7 <https://doi.org/10.1186/1471-2393-11-7>
 48. Erlindawati, Chompikul J, Isaranurug S. Factors related to the utilization of antenatal care services among pregnant women at health centers in Aceh Besar district, Nanggroe Aceh Darussalam prov-ince, Indonesia. J Public Health Dev. 2008; 6:99-108.
 49. Tabrizi FM, & Barjasteh S. Maternal hemoglobin levels during pregnancy and their association with birth weight of neonates. Iranian Journal of Pediatric Hematology and Oncology. 2015;5(4), 211

How to cite this article: Acheampong K, Appiah S, Baffour-Awuah D et al. Prevalence of anemia among pregnant women attending antenatal clinic of a selected hospital in Accra, Ghana. Int J Health Sci Res. 2018; 8(1):186-193.
