

Review Article

Relationship between Periodontal Disease and Sleep Apnoea: Current Literature & Review

Dr. Dhawal Mody¹, Dr. Vrushali Lathiya², Dr. Kamalkishor Mankar¹

¹Associate Professor, ²Senior Lecturer,
VSPM Dental College & Research Centre, Department of Periodontology, Digdoh Hills, Hingna Road,
Nagpur - 440019

Corresponding Author: Dr. Dhawal Mody

ABSTRACT

Obstructive Sleep Apnoea (OSA) is a sleep-related respiratory condition that involves a temporary cessation of breathing or complete halt in airflow despite an ongoing effort to breathe. Undiagnosed OSA is closely associated with serious health problems, including hypertension, diabetes and cardiac related disorders. Periodontitis is characterized by local bacterial infection within tooth supporting structures leading to destruction of periodontium. It is a previously unrecognized but highly prevalent and clinically relevant disease associated with OSA. It also adversely impacts cardiovascular health by increasing the production of cytokines, promoting atherosclerosis, which is also highly associated with OSA.

Keywords: sleep apnoea, periodontitis, cytokines

INTRODUCTION

Obstructive sleep apnoea (OSA) is a highly prevalent disorder affecting between 10–50% of middle-aged men. [1] OSA is characterized by recurrent upper airway obstruction associated with cyclic changes in oxyhemoglobin saturation, intermittent arousals from sleep and alterations in intrathoracic pressure. These alterations induce oxidative stress, sympathetic activation, and metabolic dysregulation. [2] OSA is currently diagnosed with an overnight sleep diagnostic test known as a poly-somnogram (PSG), which remains the gold standard of diagnosis. [3] The standard definition of an apnoeic event includes a minimum 10-second interval between breaths, with a neurologic arousal, a blood oxygen desaturation of 3% to 4% or greater, or both arousal and desaturation. [4-6]

Prolonged SF is also implicated in vascular changes similar to the enhanced

atherosclerosis and cardiovascular morbidities associated with OSA. [7,8] In children experiencing OSA, elevated systemic levels of lipopolysaccharide (LPS)-binding protein, a surrogate marker of underlying low-grade LPS endotoxemia from the gut, [9] were independently associated with BMI and with measures of OSA severity, as well as with established measures of insulin resistance. The apnoea-hypopnoea index (AHI) is used to determine the presence of obstructive sleep apnoea (OSA). An AHI score of 5 is indicative of the presence of OSA, a common disorder characterized by repeated obstructions of breathing during sleep and known to be prevalent in 24-38% of middle aged men in western countries [10-12] and in 42% of middle-aged men in Korea. [13]

Periodontal disease is a chronic inflammatory disease and its prevalence

increases with age. Risk factors for periodontal disease are age, smoking, obesity and various other socioeconomic factors. Recently it has been proved that periodontal diseases have relationship with various systemic diseases like diabetes, coronary disease, osteoporosis and low birth weight deliveries. Obstructive sleep apnoea is another one of these conditions associated with an inflammatory response. [14]

Relationship between Sleep Apnoea and Pro-inflammatory Cytokines

Recently, periodontitis has received increasing attention because it may have systemic effects, including infective endocarditis, coronary heart disease, diabetes mellitus, respiratory diseases, osteonecrosis and even malignancy. [15-19] Interleukin (IL)-1 β and IL-33 are associated with both acute and chronic inflammation. [20] Increased levels of IL-1 β were detected in saliva, gingival tissue, and gingival crevicular fluid (GCF) of patients with periodontitis. [21-23] The prominent biologic effect of IL-33 is the induction of T-helper 2 cytokines, [24] and it also possesses anti-inflammatory activity. [25]

Pro-inflammatory cytokines are also up-regulated in patients with Obstructive Sleep Apnoea (OSA). [26-28] In particular, significant elevations in serum levels of tumor necrosis factor- α (TNF- α), interleukin 1 β (IL-1 β), and interleukin 6 (IL-6) have been seen in patients with obstructive sleep apnoea. [29-33] The recent meta-analysis (MA) showed that there is an increase in levels of inflammatory markers in subjects with OSA including CRP, TNF α , IL-6, IL-8, ICAM, VCAM and Selectins. This effect is positively influenced by severity of OSA. [34]

Recent evidences highlighting the relationship between Obstructive Sleep Apnoea and Periodontitis

A preliminary study conducted by Seo WH et al. showed significant association between OSA and periodontal disease. The results showed that 17.5% of the participants had periodontitis, 46.6% had OSA and 60.0% who were diagnosed

with periodontitis had OSA. Authors identified old age, male gender, current smoking status and mouth breathing during sleep as risk factors for periodontitis. OSA was positively associated with periodontitis in subjects with 55 years of age, but not in subjects < 55 years of age. [35]

Keller et al looked at associations between obstructive sleep apnoea and chronic periodontitis in a population-based case control study. They included 7673 subjects with OSA as cases and randomly selected 21,963 subjects without a history of OSA as controls and performed conditional logistic regression analyses to explore the association between OSA and having been previously diagnosed with CP. The authors found significant difference in the prevalence of prior chronic periodontitis between cases and controls. They concluded there is associated between OSA and periodontitis. [36]

A Small Exploratory study conducted in Australia by Gunaratnam, et al reported a significant positive relationship between polysomnography-derived apnoea hypopnoea index with less than 5 events/hour and periodontitis. [37]

A case-control study conducted by Ahmad NE et al found a positive significant association between moderate or severe periodontitis. They further concluded that the patients with moderate to severe periodontitis were 4.1 times more likely to be at risk for obstructive sleep apnoea than the control patients with gingivitis or slight periodontitis (p=0.007). [38]

Latorre et al conducted a recent study to identify the association between periodontal disease and OSA in adults with different comorbidities. One hundred and ninety-nine individuals (107 women and 92 men) underwent polysomnography with a mean age of 49.9 years were recruited. They reported a significant link between Periodontitis and mild OSA and this association was more frequent in women with hypertensive cardiomyopathy. In addition, periodontitis was associated with severe OSA in men who showed any of two

comorbidities such as hypertension or hypertensive cardiomyopathy. [39]

Hikmet Gamsiz Isik et al compared periodontitis prevalence between controls and patients with OSA by assessing clinical periodontal parameters and gingival crevicular fluid (GCF) levels of interleukin (IL-1 β), tumor necrosis factor (TNF- α), and high-sensitive C-reactive protein (hs - CRP); serum hs - CRP was also sampled. They concluded higher prevalence of periodontitis and higher levels of GCF IL - 1 β and serum hs - CRP in patients with OSA. [40]

On the contrary cross sectional study conducted by Loke W et al concluded that Obstructive Sleep Apnoea (OSA) was not significantly associated with the prevalence of moderate/severe periodontitis and the periodontal parameters examined, except percentage plaque. [41]

CONCLUSION

There appears to be some evidence indicating higher levels of markers of systemic inflammation in patients with Periodontitis is associated with Obstructive Sleep Apnoea. Still this association is in very preliminary stage and further studies are needed to explore the role of these markers on the progression of Obstructive Sleep Apnoea both from the research and clinical perspective.

REFERENCES

1. Heinzer R, Vat S, Marques-Vidal P, Marti-Soler H, Andries D, Tobback N, et al. Prevalence of sleep-disordered breathing in the general population: the HypnoLaus study. *Lancet Respir Med.* 2015; 3:310–8.
2. Sánchez-de-la-Torre M, Campos-Rodriguez F, Barbé F. Obstructive sleep apnoea and cardiovascular disease. *Lancet Respir Med.* 2013; 1:61–72.
3. Epstein LJ, Kristo D, Strollo PJ Jr., et al; Adult Obstructive Sleep Apnea Task Force of the American Academy of Sleep Medicine. Clinical guideline for the evaluation, management and long-term care of obstructive sleep apnea in adults. *J Clin Sleep Med* 2009; 5:263-276.
4. [No authors listed]. Sleep-related breathing disorders in adults: Recommendations for syndrome definition and measurement techniques in clinical research. The Report of an American Academy of Sleep Medicine Task Force. *Sleep* 1999; 22:667-689.
5. Redline S, Budhiraja R, Kapur V, et al. The scoring of respiratory events in sleep: Reliability and validity. *J Clin Sleep Med* 2007; 3:169-200.
6. Iber CA, Chesson AL, Quan SF. The AASM Manual for the Scoring of Sleep and Associated Events: Rules, Terminology, and Technical Specifications. Westchester, IL: American Academy of Sleep Medicine; 2007.
7. A Carreras, S.X. Zhang, E. Peris, et al. Chronic sleep fragmentation induces endothelial dysfunction and structural vascular changes in mice. *Sleep* 2014; 37(11):1817-1824.
8. B Mokhlesi, S.A. Ham, D. Gozal The effect of sex and age on the comorbidity burden of OSA: an observational analysis from a large nationwide US health claims database. *Eur Respir J* 2016; 47(4):1162-1169
9. L. Kheirandish-Gozal, E. Peris, Y. Wang, et al. Lipopolysaccharide-binding protein plasma levels in children: effects of obstructive sleep apnea and obesity. *J Clin Endocrinol Metab* 2014; 99:656-663.
10. Olson LG, King MT, Hensley MJ, Saunders NA. A community study of snoring and sleep-disordered breathing Prevalence. *Am J Respir Crit Care Med* 1995; 152:711–716.
11. Kripke DF, Ancoli-Israel S, Klauber MR, Wingard DL, Mason WJ, Mullaney DJ. Prevalence of sleep-disordered breathing in ages 40-64 years: a population-based survey. *Sleep* 1997; 20:65–76.
12. Bearpark H, Elliott L, Grunstein R et al. Snoring and sleep apnea. A population study in Australian men. *Am J Respir Crit Care Med* 1995; 151:1459–1465.
13. Kim J, In K, You S et al. Prevalence of sleep-disordered breathing in middle-aged Korean men and women. *Am J Respir Crit Care Med* 2004; 170:1108–1113.
14. Ryan S, Taylor CT, McNicholas WT. Selective activation of inflammatory pathways by intermittent hypoxia in obstructive sleep apnoea syndrome. *Circulation*, 2005; 112(17):2660-2667.
15. Azarpazhooh A, Leake JL. Systematic review of the association between

- respiratory diseases and oral health. *J Periodontol* 2006; 77:1465–1482.
16. Ezzo PJ, Cutler CW. Microorganisms as risk indicators for periodontal disease. *Periodontol* 2000 2003; 32:24–35.
 17. Mustapha IZ, Debrey S, Oladubu M, Ugarte R. Markers of systemic bacterial exposure in periodontal disease and cardiovascular disease risk: a systematic review and meta-analysis. *J Periodontol* 2007; 78:2289–2302.
 18. Geurs NC. Osteoporosis and periodontal disease. *Periodontol* 2000 2007; 44:29–43.
 19. Ritchie CS. Obesity and periodontal disease. *Periodontol* 2000 2007; 44:154–163.
 20. Meyer MS, Joshupura K, Giovannucci E, Michaud DS. A review of the relationship between tooth loss, periodontal disease, and cancer. *Cancer Causes Control* 2008; 19:895–907.
 21. Van de Veerdonk FL, Netea MG. New Insights in the immunobiology of IL-1 family members. *Front Immunol* 2013; 4:167.
 22. Gu'rsoy UK, Ko'no'nen E, Uitto VJ, et al. Salivary interleukin-1beta concentration and the presence of multiple pathogens in periodontitis. *J Clin Periodontol* 2009; 36:922-927.
 23. Howells GL. Cytokine networks in destructive periodontal disease. *Oral Dis* 1995; 1:266-270.
 24. Louten J, Rankin AL, Li Y, et al. Endogenous IL-33 enhances Th2 cytokine production and T-cell responses during allergic airway inflammation. *Int Immunol* 2011; 23:307-315.
 25. Cohen I, Rider P, Carmi Y, et al. Differential release of chromatin-bound IL-1alpha discriminates between necrotic and apoptotic cell death by the ability to induce sterile inflammation. *Proc Natl Acad Sci USA* 2010; 107:2574-2579.
 26. Ursavaş A, Karadağ M, Rodoplu E, Yilmaztepe A, Oral HB, Gözü RO. Circulating ICAM-1 and VCAM-1 levels in patients with obstructive sleep apnea syndrome. *Respiration* 2007; 74:525-32.
 27. Von Känel R, Loredó JS, Ancoli-Israel S, Mills PJ, Natarajan L, Dimsdale JE. Association between polysomnographic measures of disrupted sleep and prothrombotic factors. *Chest* 2007; 131:733-9.
 28. Punjabi NM, Beamer BA. C-reactive protein is associated with sleep disordered breathing independent of adiposity. *Sleep* 2007; 30:29-34.
 29. De la Pena Bravo M, Serpero LD, Barcelo A, et al. Inflammatory proteins in patients with obstructive sleep apnoea with and without daytime sleepiness. *Sleep Breath* 2007; 11:177-85.
 30. Minoguchi K, Tazaki T, Yokoe T, et al. Elevated production of tumor necrosis factor- α by monocytes in patients with obstructive sleep apnea syndrome. *Chest* 2004; 126:1473-9.
 31. Ciftci TU, Korturk O, Bukan N, et al. The relationship between serum cytokine levels with obesity and obstructive sleep apnoea syndrome. *Cytokine* 2004; 28:87-91.
 32. Vgontzas AN, Papanicolaou DA, Bixler EO, Kales A, Tyson K, Chrousos GP. Elevation of plasma cytokines in disorders of excessive daytime sleepiness: role of sleep disturbance and obesity. *J Clin Endocrinol Metab* 1997; 82:1313-6.
 33. Vgontzas AN, Papanicolaou DA, Bixler EO, et al. Sleep apnea and daytime sleepiness and fatigue: relation to visceral obesity, insulin resistance, and hypercytokinemia. *J Endocrinol Metab* 2000; 85:1151-8.
 34. Nadeem R, Molnar J, Madbouly EM, Nida M, Aggarwal S, Sajid H, Naseem J; Loomba R. Serum inflammatory markers in obstructive sleep apnoea: a meta-analysis. *J Clin Sleep Med* 2013;9(10):1003-1012.
 35. Seo WH, Cho ER, Thomas RJ, An S-Y, Ryu J, Kim H, Shin C. The association between periodontitis and obstructive sleep apnea: a preliminary study. *J Periodont Res* 2013; 48: 500–506.
 36. Keller JJ, Wu CC, Chen YH, Lin HC. Association between obstructive sleep apnea and chronic periodontitis: a population-based study. *J Clin Periodontol*, 2013; 40(2):111-117.
 37. Gunaratnam K, Taylor B, Curtis B, Cistulli P. Obstructive sleep apnoea and periodontitis: A novel association? *Sleep Breath*. 2009; 13(3):233–239.
 38. Ahmad NE, Sanders AE, Sheats R, Brame J, Essick GK. Obstructive sleep apnoea in association with periodontitis: a case-control study. *J Dental Hyg*, 2013; 87(4):188-199.
 39. Catalina Latorre, Francina Escobar, Juliana Velosa, Daniela Rubiano, Patricia Hidalgo-Martinez, Liliana Otero. Association between obstructive sleep apnea and comorbidities with periodontal disease in

- adults. *J Indian Soc Periodontol.* 2018 May-Jun; 22(3): 215–220.
40. Hikmet Gamsiz Isik , Esen Kiyani, Zuleyha Bingol , Ulku Baser, Evin Ademoglu, Funda Yalcin,. Does Obstructive Sleep Apnea Increase the Risk for Periodontal Disease? A Case Control Study. *J Periodontol* 2017; 88:443-449.
41. Weiqiang Loke, Thomas Girvan, Paul Ingmundson, Ronald Verrett, John Schoolfield, Brian L. Mealey. Investigating the Association Between Obstructive Sleep Apnoea and Periodontitis. *J Periodontol* 2015;86:232-243.

How to cite this article: Mody D, Lathiya V, Mankar K. Relationship between periodontal disease and sleep apnoea: current literature & review. *Int J Health Sci Res.* 2019; 9(1):246-250.
